

## Highlights

- Subjects were selectively bred High Runner (HR) and non-selected Control (C) mice.
- Juveniles were housed with/without wheels and given standard or Western diet (WD).
- After 8 weeks washout, early-life WD increased adult wheel running only in HR mice.
- Early exercise increased muscle and brain mass, and reduced anxiety-like behavior.
- Early-life WD increased adult preference for WD but not for sucrose.

Effects of early-life exposure to Western diet and voluntary exercise on adult activity levels, exercise physiology, and associated traits in selectively bred High Runner mice

Marcell D. Cadney<sup>a</sup>, Layla Hiramatsu<sup>a</sup>, Zoe Thompson<sup>b</sup>, Meng Zhao<sup>a</sup>, Jarren C. Kay<sup>a</sup>, Jennifer M. Singleton<sup>a</sup>, Ralph Lacerda de Albuquerque<sup>a</sup>, Margaret P. Schmill<sup>b</sup>, Wendy Saltzman<sup>a</sup>, Theodore Garland, Jr.<sup>a, \*</sup>

<sup>a</sup> Department of Evolution, Ecology, and Organismal Biology, University of California, Riverside, CA 92521, USA

<sup>b</sup> Neuroscience Graduate Program, University of California, Riverside, CA 92521, USA

\* Corresponding author.

E-mail address: tgarland@ucr.edu (T. Garland).

## Abstract

Exercise behavior is under partial genetic control, but it is also affected by numerous environmental factors, potentially including early-life experiences whose effects persist into adulthood. We studied genetic and early-life environmental effects on wheel-running behavior in a mouse model that includes four replicate high runner (HR) lines selectively bred for increased voluntary wheel running as young adults and four non-selected control (C) lines. In a full factorial design, mice from each line were granted wheel access or not and administered either standard or Western diet (WD) from weaning (3 weeks old) to 6 weeks of age (sexual maturity). In addition to acute effects, after a washout period of 8 weeks (~6 human years) in which all mice had standard diet and no wheel access, we found both beneficial and detrimental effects of these early-life exposures. During the first week of treatments, WD increased distance run by 29% in C mice and 48% in HR mice (significant Diet  $\times$  Linetype interaction), but diet effects disappeared by the third week. Across the three weeks of juvenile treatment, WD significantly increased fat mass (with lean mass as a covariate). Tested as adults, early-life exercise increased wheel running of C mice but not HR mice in the first week. Early-life exercise also reduced adult anxiety-like behavior and increased adult fasted blood glucose levels, triceps surae mass, subdermal fat pad mass, and brain mass, but decreased heart ventricle mass. Using fat mass as a covariate, early-life exercise treatment increased adult leptin concentration. In contrast, early-life WD increased adult wheel running of HR mice but not C mice. Early-life WD also increased adult lean mass and adult preference for Western diet in all groups. Surprisingly, early-life treatment had no significant effect on adult body fat or maximal aerobic capacity ( $VO_{2max}$ ). No previous study has tested for combined or interactive effects of early-life WD and exercise. Our results demonstrate that both factors can have long-lasting effects on adult voluntary exercise and related phenotypes, and that these effects are modulated by genetic background. Overall, the long-lasting effects of early-life exercise were more pervasive than those of WD, suggesting critical opportunities for health intervention in childhood habits, as well as possible threats from modern challenges. These results may be relevant for understanding potential effects of activity reductions and dietary changes associated with the obesity epidemic and COVID-19 pandemic.

**Keywords:** Developmental programming; Early-life effects; Exercise; Reward; Western diet; Wheel running

## Highlights

- Subjects were selectively bred High Runner (HR) and non-selected Control (C) mice.
- Juveniles were housed with/without wheels and given standard or Western diet (WD).
- After 8 weeks washout, early-life WD increased adult wheel running only in HR mice.
- Early exercise increased muscle and brain mass, and reduced anxiety-like behavior.
- Early-life WD increased adult preference for WD but not for sucrose.

## 1. Introduction

Human obesity and most of its negative health consequences result from complex interactions among diet, physical activity, environmental exposures, sex, genetic predisposition, and sociocultural factors (Eisenmann 2004, 2006; Ross and Desai 2005; Adam and Epel 2007; Papas et al. 2007; Choh et al. 2009; Coccurello et al. 2009; McAllister et al. 2009; Sisson et al. 2009; Stein et al. 2009; Kirk et al. 2010; North et al. 2010; Graff et al. 2011; Patterson and Abizaid 2013; Yang and Huffman 2013; Rohde et al. 2019). Of these, declining physical activity and "Western" diets (high in fat & simple sugars) are seen as key contributors to increasing rates of obesity and its comorbidities in all age groups across multiple populations (Cornier et al. 2008; Westerterp and Speakman 2008; Steinberger et al. 2009; Flegal et al. 2010; Halpern et al. 2010). Conversely, a "good" diet (e.g., relatively low in sugar and fat [but see Esposito et al. (2010); O'Neill and Raggi (2020)]) and higher activity levels promote physical fitness and cardiovascular health while lowering the risk for obesity, Type 2 diabetes, and their comorbidities (Blair and Morris 2009; Hills et al. 2011; McGuire and Ross 2011; Janssen et al. 2013; Lee et al. 2017). Importantly, not only voluntary exercise (VE) but also incidental or spontaneous physical activity (SPA) has beneficial effects in both animal models and humans (e.g., Levine 2007; Levine et al. 2008; Novak et al. 2009; Garland, Jr. et al. 2011; McGuire and Ross 2011; Teske et al. 2014; Lightfoot et al. 2018).

Like obesity itself, human physical activity and diet choices are products of both genes and numerous environmental effects acting throughout an individual's development (Trost et al. 2002; Lightfoot et al. 2018). Many key environmental factors are experienced early in life, including *in utero* (Garland, Jr. et al. 2017). Such ontogenetic factors can strongly contribute to the development of various unhealthy adult behaviors, habits, diseases, and disorders (Dishman et al. 1985; Sallis and Hovell 1990; Twisk 2001; Barker 2003; de Boo and Harding 2006; Nathanielsz 2006; Hanson and Gluckman 2008; Gardner and Rhodes 2009; Langley-Evans 2009; Metges 2009; Ozanne and Siddle 2011; Blondin et al. 2016). Unfortunately, interventions to improve diet or increase physical activity often have little impact on children's activity (Wilkin 2011; Roberto et al. 2015; but see Gomersall et al. 2016; Reilly 2011), and have not halted the increasing prevalence of obesity in children (Harris et al. 2009).

Human studies suggest that experiences occurring from conception to sexual maturity can alter adult levels of voluntary exercise and/or SPA. For example, adults who were exposed to the Dutch Famine *in utero* have increased adiposity and more atherogenic lipid profiles that may be related to decreased physical activity (Lussana et al. 2008; Stein et al. 2009). However, mechanisms by which early-life experiences affect physical activity are poorly understood (Dishman et al. 1985; Twisk 2001; Jung et al. 2006; Lussana et al. 2008; Andersen et al. 2009; Downing et al. 2011; Koenen et al. 2011; Li et al. 2013; Baker et al. 2015), in part because existing human studies are mostly correlational, cross-sectional (Trost et al. 2002), or of questionable methodological quality (e.g., use of questionnaires to gauge physical activity: Hallal et al. 2006; Stein et al. 2009; Koenen et al. 2011).

Numerous rodent studies establish that *in utero* and other early-life exposures to calorie-restricted, low-protein, high-fat, or high-carbohydrate diets can affect obesity-related measures in adults, such as body composition, appetite, dietary preferences, reward signaling, and the microbiome (Frazier et al. 2008; Teegarden et al. 2009; Vucetic et al. 2010, 2012; Ozanne and Siddle 2011; Sun et al. 2013; Desai et al. 2014; McNamara et al. 2021). Far fewer studies address maternal or early-life effects on adult voluntary exercise or SPA (Breier et al. 2006; Dai et al. 2012; Sun et al. 2013; Hiramatsu et al. 2017; see also Donovan et al. 2013 on sheep).

Here, we aimed to identify possible long-lasting effects of juvenile exposure to Western diet (WD) and exercise. We used mice from a well-characterized experimental evolution animal model: four replicate High Runner (HR) lines that were selectively bred for 76 generations for increased voluntary wheel-running behavior on days 5 and 6 of a 6-day wheel exposure as young adults and four non-selected control (C) lines (Swallow et al. 1998; Careau et al. 2013). On average, mice from the HR lines run ~3-fold greater distance per day compared with the C lines, for both sexes (Swallow et al. 1998). A number of other phenotypes have evolved in association with wheel-running behavior in the HR lines, including greater SPA when wheels are not provided, increased running endurance during forced treadmill exercise, increased maximal aerobic capacity during forced exercise (VO<sub>2</sub>max), reduced body fat, larger hearts, altered dopamine signaling in the brain, altered endocannabinoid signaling, higher baseline circulating corticosterone levels, and lower circulating leptin levels (Rhodes et al. 2005; Rhodes and Kawecki 2009;

Swallow et al. 2009; Garland, Jr. et al. 2011, 2016; Keeney et al. 2012; Wallace and Garland, Jr. 2016; Thompson et al. 2017; Kay et al. 2019).

Previously, we found that access to Western diet beginning at weaning and continuing through adulthood had a large stimulatory effect (~50–75%) on daily wheel-running distance of both juvenile and adult male HR mice, with no statistically significant effect on C mice (Meek et al. 2010; only males tested). These dramatic results for voluntary exercise were unprecedented for rodents, although in adult rats a high-fat diet can increase VO<sub>2</sub>max and endurance capacity (Simi et al. 1991). A subsequent study of HR and C mice found that early-life wheel access for just 3 weeks, starting at weaning, increased adult wheel running in males of both linetypes, with no effect on SPA and contrasting effects on adult circulating leptin levels in HR and C mice (Acosta et al. 2015). All early-life-exercise mice had significantly reduced adult body mass and a trend for reduced visceral fat pad mass (Acosta et al. 2015). Taken together, these previous results demonstrate that early-life effects on adult physical activity can be genotype-dependent.

In the present study, we addressed three specific questions. (1) Do WD and wheel access given to juveniles have acute/immediate effects on juvenile physical activity, caloric intake, and body composition? (2) Do WD and wheel access given to juveniles have long-lasting effects on adult physical activity and related phenotypes? (3) Do early-life diet or exercise interact with genetic backgrounds to influence adult phenotypes? From what has been reported in similar previous studies (Meek et al. 2010; Acosta et al. 2015), we expected that WD and/or exercise would have substantial effects when applied during the juvenile period, and we hypothesized that those effects would carry over into adulthood, potentially affecting locomotor behavior (Garland, Jr. et al. 2017). To address these questions, we used HR and C male mice from all replicate lines of generation 76 of the HR animal model. During the juvenile period (3-6 weeks post birth), we administered either standard diet or WD and altered voluntary exercise by granting wheel access or not. We tested juveniles at 6 weeks for SPA, caloric intake, and body composition. After a washout period of eight weeks (equivalent to ~6 human years), we then tested adults for VE, SPA, and related traits.

## 2. Materials and Methods

### 2.1. Experimental mice

Starting in 1993, four replicate lines of house mice were bred in an ongoing selection experiment for high voluntary wheel running (HR lines), based on the number of wheel revolutions on days five and six of six days of access to Wahman-type activity wheels (1.12-meter circumference) as young adults (Swallow et al. 1998). The experiment began with a population of 224 mice from the outbred Hsd:ICR strain, which was randomly mated for two generations before being randomly partitioned into eight lines. Four of these were bred randomly as Control (C) lines to the four HR lines. The current experiment used a subset of mice from generation 76. All experiments followed University of California Riverside IACUC guidelines.

### 2.2. Early-life diet and exercise manipulation

In this experiment, 196 male mice (half HR and half C) were weaned individually into standard cages at 3 weeks of age and placed in one of four treatment groups for 3 weeks until sexual maturity at 6 weeks of age (Fig. 1). Only males were studied due to resource constraint (e.g., the number of wheels available) and to avoid complications caused by possible estrus-cycle effects on some of the outcome variables. Half of the mice from each line were given “Western” diet (WD: Harlan Teklad TD.88137, 42% kJ from fat [anhydrous milk], 42.7% kJ from carbohydrates [sucrose and cornstarch], 15.2% kJ from protein [casein], 34.1% added sucrose by weight) and the other half were given standard diet (SD: Teklad Rodent Diet W-8604, 14% kJ from fat, 54% kJ from carbohydrates, and 32% kJ from protein, no added sugars [less than ~9% naturally occurring sugars by weight, mostly from grains]). According to Teklad/Envigo, the protein sources are very different for these two diets, so the amino acid profiles may be different as well, but both should meet all of the estimated requirements for amino acids for rodents (Derek Martin in email of 8 Feb. 2021). See Meek et al. (2010) for more nutritional information on the Western and standard mouse chow.

Within each diet group, half of the animals from each line occupied cages attached to activity wheels (see above). Body mass, food consumption, and body composition



1  
2  
3  
4 were measured every seven days after weaning until mice were 6 weeks old.  
5 Photoperiod was 12:12, with lights on at 07:00 h.  
6

### 7 *2.3. Adult testing*

8  
9 Beginning at 6 weeks of age, all mice were individually housed with SD and  
10 without wheel access for an additional 8 weeks. At 14 weeks of age, testing of VO<sub>2</sub>max,  
11 open-field behavior, WD and sucrose preference began. At 18 weeks of age, blood lipid  
12 levels were measured, and all adult mice were given access to wheels for 2 weeks to  
13 measure VE. Adult SPA was again measured for the subset of mice that did not have  
14 access to wheels as juveniles. Mice were euthanized and tissues collected at  
15 approximately 20 weeks of age.  
16  
17

18 For each day of wheel testing (3 weeks as juveniles and 2 weeks as adults), we  
19 recorded revolutions in each 1-minute interval over a period of 23 h. This allowed  
20 computation of the total distance run, the number of 1-minute intervals with at least one  
21 revolution (a measure of the daily running duration), and the mean running speed  
22 (distance/intervals), as well as the maximum speed in any 1-minute interval (Copes et al.  
23 2015). We also computed weekly means for these measures. Wheel freeness (an  
24 inverse measure of how difficult it is to turn the wheel) was used as a covariate in all  
25 analyses of wheel running (e.g., Copes et al. 2015).  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38

#### 39 *2.3.1. Home-cage activity*

40  
41 SPA was measured as home-cage activity. Mice were housed in home cages  
42 fitted with a passive infrared sensor (Talon TL-Xpress-A; Crow Electronics, Fort Lee, New  
43 Jersey, USA), protected within wire mesh, as in previous studies (Acosta et al. 2015;  
44 Copes et al. 2015). The sensors were connected to a computer with custom activity-  
45 recording software (developed by M. A. Chappell) via a digital I/O board (ICS 2313; ICS  
46 Electronics, Pleasanton, CA, USA). The sensors recorded activity three times per  
47 second and a mean value between 0 (no movement detected) and 1 (movement  
48 detected) was calculated for each minute over 23 hour-periods of measurement. All  
49 analyses of SPA data used a measure of sensor sensitivity as a covariate (Acosta et al.  
50 2015; Copes et al. 2015). Juvenile and adult SPA could only be measured in mice that  
51 were not given wheel access during the juvenile treatment stage.  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

### 2.3.2. *Body composition*

Whole-animal fat, lean, free water, and total water masses were measured by restraining each mouse within a translucent tube before insertion into an EchoMRI-100 (Echo Medical Systems, Houston, TX, USA) for scanning. This procedure lasted approximately 1-2 min per mouse and did not require sedation or anesthesia. Body composition was analyzed at weaning, after each of the three weeks of juvenile exposure, and again at 9 weeks of age, prior to adult wheel testing.

### 2.3.3. *Caloric intake*

Food consumption was measured by the change in food hopper weight each week during early-life treatment (when both SD and WD were used) and during adult wheel testing (when all mice received SD), taking care to note wasted or shredded food (Koteja et al. 2003). Weekly food consumption was converted from grams to caloric intake, taking the caloric content of SD and WD into account (14 and 19 kJ per gram, respectively; Meek et al. 2010).

### 2.3.4. *Maximal aerobic capacity*

To measure  $\text{VO}_2\text{max}$ , mice were subjected to forced exercise within a 900 mL enclosed mouse wheel approximately 15 cm in diameter (Dlugosz et al. 2009; Claghorn et al. 2017). Each mouse was run for approximately 5 min. while researchers manually accelerated the wheel. Duplicate trials were conducted, allowing a day of rest between each trial. Air was pumped into the enclosed metabolic chamber at a rate of 2000 mL per min, and the concentration of  $\text{O}_2$  in excurrent air was measured by an oxygen analyzer (S-3A Applied Electrochemistry, Inc., Sunnyvale, CA). We then calculated oxygen consumption as the difference in  $\text{O}_2$  concentration between reference air (pumped through the wheel before the mouse was placed inside) and sampled air (while the mouse was inside). Following previous studies (Claghorn et al. 2017; Hiramatsu et al. 2017),  $\text{VO}_2\text{max}$  was taken as the highest minute of oxygen consumption during either trial, as calculated with LabHelper software (Warthog Systems, [www.warthog.ucr.edu](http://www.warthog.ucr.edu)). Trials that did not evidence a plateau in  $\text{O}_2$  with increasing wheel speed were excluded.

Researchers subjectively assessed mouse cooperation during each trial and tiredness after each trial to assess the quality of the trial. Cooperation was scored

between 1, least cooperative (the mouse made repeated attempts to breathe through the chamber aperture and/or resisted moving in the direction of motion) and 5, most cooperative (the mouse was least interested in the aperture and focused mostly on running with the direction of rotation). Tiredness was scored between 1, least exhausted after the trial, to 3, most exhausted. Tiredness (1-3 scale) was scored based on long it took for the mouse to begin moving about the respiration chamber after the end of forced exercise (1 = 1 s or less; 3 = 5 s or more). The identity of the investigator scoring each trial was recorded, so that potential differences in scoring could be accounted for. Trials were excluded from analyses if cooperation scores were less than 3 or tiredness scores were less than 2, as values this low likely correspond to trials in which VO<sub>2</sub>max would not be attained.

#### *2.3.5. Preference for Western diet and sucrose solution*

At approximately 17 weeks of age, adult mice were presented in their home cage with partitioned food hoppers containing WD and SD for 24 h (13:00–13:00 h). Consumption of each diet was measured as the change in hopper weight across the test period, with due allowance for spillage and wasting (Koteja et al. 2003). The next day, mice were given a similar choice between water and sucrose-water (13:00–13:00 h) (10.5% sucrose; Fisher Scientific Certified ACS Grade). WD preference was calculated by caloric conversion: WD kJ / (WD kJ + SD kJ) (Meek et al. 2010; Acosta et al. 2017). Sucrose preference was calculated similarly as volume of sucrose-water / (volume of sucrose-water + volume of water).

#### *2.3.6. Open-field behavior*

Between 15 and 16 weeks of age, mice were tested for open-field behavior (Fuss et al. 2009) on reversed photoperiod (at approximately 07:00–18:00 h). Each mouse was placed in the center of an arena (1 m<sup>2</sup>) surrounded by 0.5 m black Trovicel walls under low-light conditions (Bronikowski et al. 2001). The mouse was recorded for 5 min with a Logitech HD C525 Webcam placed above the arena. Videos were assessed with TopScan LITE software (Clever Sys Inc., Reston, VA). The amount of urine (scored as 0, 1, 2) and the number of fecal pellets left by the mouse during the test were recorded. The arena was cleaned with warm, soapy water and dried between each trial. Analyses

of time spent near the arena walls were done on each min of the trials, using time of day as a covariate.

### *2.3.7. Blood lipid profile*

At 17 weeks of age, food was removed at 07:00 h and blood samples were taken from the submandibular vein using sterile, single-use lancets under isoflurane anesthesia at 13:00–19:00 h (Golde et al. 2005). Total cholesterol (TC), high-density lipoprotein (HDL) cholesterol, triglycerides, and glucose (GLU) in whole blood, as well as TC/HDL (total cholesterol/HDL cholesterol) ratio and estimates of low-density lipoprotein (LDL) and non-HDL cholesterol were measured and calculated by an Abbott Cholestech LDX™ Analyzer. Time of day and bleed delay time (elapsed time from disturbing the mouse to collecting a blood sample) were used as covariates in statistical analyses.

### *2.3.8. Dissections and plasma hormone concentrations*

Mice were removed from wheel access for one day then euthanized via decapitation. Trunk blood was immediately collected into heparinized containers, and organs (brain, heart ventricles, spleen, liver, triceps surae muscles, and fat pads) were dissected and weighed to 0.0001 g. Plasma samples were prepared and preserved as instructed by the assay kits described below.

Terminal plasma leptin was measured in duplicate using a Crystal Chem enzyme-linked immunosorbent assay (ELISA) kit (Mouse Leptin Assay Catalog #90030) without dilution. Absorbances were read at 450 nm in an EPOCH2 microplate reader using GEN5 2.07 reading software (microplate and reading software: BioTek Instruments, Winooski, VT, USA). Terminal plasma adiponectin was measured in duplicate with an AssayPro ELISA kit (Mouse Adiponectin ACRP30 Catalog #EMA 2500-1), using a 400-fold dilution. Prior to dissection, plasma was collected from the caudal vein at 17 weeks of age (Fig. 1) specifically to measure corticosterone concentration in duplicate with an Arbor Assays ELISA kit (Corticosterone EIA kit Catalog #K014-H1), using a 100-fold dilution. All assay plates were read in triplicate.

### 2.3.9. Statistical analysis

Data were analyzed as mixed models in SAS 9.1.3 (SAS Institute, Cary, NC, USA) Procedure Mixed, with REML estimation and Type III Tests of Fixed Effects. Depending on the trait analyzed, body mass, age, wheel freeness, and home-cage sensor sensitivity were used as covariates. Linetype (HR vs. C), diet, and wheel access were fixed effects, while replicate lines were a random effect nested within linetype. Effects of linetype, diet, and wheel access, as well as their interactions, were tested with 1 and 6 degrees of freedom. Modeling the aforementioned random effects occasionally resulted in covariance parameter estimates of zero for some of the interactive random terms. In these cases, the interactive random terms were removed, starting with higher-order terms. The line within linetype random term was never removed. Once analyses returned non-zero covariance parameter estimates, outliers were then removed when the standardized residual exceeded  $\sim 3$ .

An additional fixed effect, termed mini-muscle, was included in all statistical analyses. The original base population of the outbred strain had a mini-muscle phenotype, which is caused by a Mendelian recessive allele that is a novel intronic single nucleotide polymorphism in the *Myosin heavy polypeptide 4* gene (Kelly et al. 2013). The allele started at a frequency of  $\sim 7\%$  in the base population and was under positive selection in the HR lines, eventually going to fixation in one HR line while remaining polymorphic in another (Garland, Jr. et al. 2002; Syme et al. 2005; Hannon et al. 2008; Hillis et al. 2020). The mini-muscle phenotype is characterized by a 50% reduction in the triceps surae muscle and total hindlimb muscle mass – due largely to reduced type IIb muscle fibers (Guderley et al. 2006; Talmadge et al. 2014). The effect of mini-muscle phenotype is reported in tables of statistical results but, for simplicity and as it is not a primary focus of the present study, not mentioned in the Results. In Discussion, we summarize results of the mini-muscle phenotype.

Leptin and adiponectin concentrations were analyzed with covariates of fat mass (as measured by MRI). To remove outliers from plasma hormone data, we tested for variation due to multiple reads of the assay plates and variation between duplicates. SAS Procedure Mixed was used with Mouse ID as the independent variable and sample duplicate nested within Mouse ID and read triplicate nested within sample duplicate as random effects. Corticosterone samples with bleed delay time of 3 minutes or more were

excluded from analysis ( $n = 10$ ). As described above, outliers were removed when the standardized residual exceeded  $\sim 3$ .

To investigate any possible effects of litter sex ratio at weaning, we used sex ratio as a covariate in many relevant preliminary tests (e.g., juvenile and adult wheel running distance, body mass and composition, maximal aerobic capacity, etc.). The effect of sex ratio was never statistically significant (results not shown), so final models did not include it as a covariate.

### 3. Results

Juvenile exposure to a Western diet and/or voluntary exercise had numerous effects on juvenile and adult wheel-running behavior, body mass and fat composition, and other behavioral and physiological traits, some in an interactive genotype-by-environment manner.

#### *3.1. Juvenile wheel running and spontaneous physical activity (during treatment)*

During the first week of treatment, WD increased the distance (i.e., number of revolutions) run by 29% in C mice and by 48% in HR mice (Fig. 2A; diet  $p < 0.0001$ ; diet  $\times$  linetype  $p = 0.0072$ ). During the second week, mice on WD ran 25% more revolutions than those on SD (diet  $p = 0.0188$ ) and HR mice ran 160% more than C mice (linetype  $p = 0.0320$ ), with no interaction. Diet effects disappeared completely by the third week, and HR mice ran more than 3-fold more revolutions per day than C mice ( $p = 0.0006$ ). See Fig. 2B–D and Supplemental Table 1 for results on duration, average speed, and maximum speed.

Home-cage activity was only recorded for mice housed without wheels, and only during the third week of exposures. Diet did not affect juvenile SPA ( $p = 0.5734$ ); however, HR mice were significantly more active in the home-cage than C mice ( $p = 0.0036$ ) (Supplemental Table 1).

#### *3.2. Adult wheel running and spontaneous physical activity*

In the first week of adult testing, HR mice ran 3.5-fold more distance than C mice (Fig. 3;  $p = 0.0008$ ). Early-life exercise increased wheel running distance among C mice,

but not HR mice (exercise  $\times$  linetype  $p = 0.0512$ ). Early-life WD increased distance for HR mice, but not for C mice (diet  $\times$  linetype  $p = 0.0091$ ). In the second week, HR mice ran 3.3-fold more than C mice ( $p < 0.0001$ ; Supplemental Table 1) and early-life WD exposure significantly increased average daily running distance of HR mice ( $p = 0.0186$ ), but not C ( $p = 0.8914$ ; diet  $\times$  linetype interaction  $p = 0.0534$ ).

During the first week, daily running duration, mean speed, and maximum speed averaged greater for HR mice (all  $p \leq 0.0251$ ). However, significant diet  $\times$  linetype interactions occurred ( $p = 0.0146$ ,  $0.0414$ , and  $0.0761$ ; Fig. 3B, C, and D, respectively), whereby early-life WD decreased these measures among C mice but increased them among HR. Early-life wheel access increased mean and maximum speeds, but not running duration (all  $p \leq 0.0086$ ). See Supplemental Table 1 for results from the second week of adult wheel running.

As noted above, only mice that did not have wheel access during the juvenile period were measured. SPA of adult mice housed with wheel access was unaffected by juvenile diet ( $p = 0.4535$ ) and did not differ between HR and C mice ( $p = 0.2218$ ), with no diet  $\times$  linetype interaction ( $p = 0.3496$  for first week; results for week two in Supplemental Table 1).

### 3.3. Body mass and composition

#### 3.3.1. Body mass

Body mass increased throughout the experiment for all four experimental groups for both HR and C lines, until finally declining during the two weeks of adult wheel testing for all groups (Fig. 4A, B). The change in body mass from the end of juvenile exposures (9 weeks of age) to the start of adult wheel testing (18 weeks of age) was not affected by juvenile treatments, with or without body mass at 9 weeks of age as a covariate. The change in body mass across the two weeks of adult wheel testing was influenced by a three-way interaction of juvenile exercise, juvenile diet, and linetype ( $p = 0.0059$ ). Mice from the HR lines were significantly smaller than C mice at all time points, including at weaning (all  $p \leq 0.0020$ ).

#### 3.3.2. Lean mass

Body lean mass increased throughout the experiment (Fig. 4C, D). Adult lean mass did not change significantly across two weeks of wheel testing. Early-life WD



increased lean mass a week after treatment ended ( $p = 0.0383$ ), and this effect persisted through washout into adulthood ( $p = 0.0316$ ). Access to wheels during the juvenile period increased lean mass after the third week of treatments ( $p = 0.0104$ ). HR mice were consistently leaner than C mice during early life and into adulthood (all  $p < 0.05$ , Supplemental Table 1).

### 3.3.3. Fat mass

During each week of juvenile treatment, WD significantly increased body fat in all groups (Fig. 4E, F; all three  $p \leq 0.0001$ , with lean mass as a covariate), with C mice on WD having the sharpest rise (compared to HR mice on WD) in body fat after 3 weeks of juvenile treatment (diet  $\times$  linetype  $p = 0.0099$ ). The apparent increase in relative body fat between 9 and 18 weeks of age was not statistically significant for any group, with no significant effect of linetype or early-life treatment. After two weeks of adult wheel testing, relative body fat was significantly reduced in all groups (all  $p \leq 0.0138$ ). At the end of adult wheel testing, relative fat mass had a three-way interaction (exercise  $\times$  diet  $\times$  linetype  $p = 0.0224$ ).

### 3.4. Caloric intake

During the first week of early-life treatment, WD increased mean daily caloric intake by ~21% (Fig. 5; diet  $p < 0.0001$ ). Among SD mice, those with wheels consumed significantly more calories than the sedentary group; however, for WD mice, those with wheel access consumed significantly fewer calories than the sedentary group (exercise  $\times$  diet  $p < 0.0001$ ). During the second week of treatment, both exercise ( $p = 0.0024$ ) and diet ( $p = 0.0009$ ) increased caloric intake, with no interactions. During the third week of treatment, mice with wheels consumed more ( $p < 0.0001$ ), and the effect was somewhat greater for HR mice (diet  $\times$  linetype  $p = 0.0881$ ).

In the first week of adult testing, the early-life treatments did not affect caloric intake. However, in the second week of adult testing caloric intake was decreased by early-life exercise, and the effect was greater among HR mice (Fig. 6; exercise  $\times$  linetype  $p = 0.0126$ ). HR mice consumed more calories than C mice during both the first (Fig. 6;  $p = 0.0276$ ) and second ( $p = 0.0178$ ) weeks of adult wheel testing.



### 3.5. Maximum aerobic capacity

With body mass as a covariate, HR mice had greater VO<sub>2</sub>max than C mice (Fig. 7;  $p = 0.0012$ ). Neither early-life exercise, nor diet had any effect on VO<sub>2</sub>max.

### 3.6. Preference for Western diet and sucrose-water

All experimental groups preferred WD to SD (see Supplemental Table 1). Early-life WD increased adult preference for Western diet in all groups, and a three-way interaction indicated that the magnitude of this effect depended on linetype and early-life exercise (see Supplemental Table 1;  $p = 0.0578$ ). All groups preferred sucrose-water to regular water, but this preference was not affected by linetype or either early-life treatment.

### 3.7. Open-field behavior

Early-life exercise reduced the time spent near the arena walls during each of the five minutes of the open-field test for all groups (all  $p < 0.05$ ). Summing across all five minutes also showed a 3-way interaction (exercise  $\times$  diet  $\times$  linetype  $P = 0.0432$ ), indicating that the magnitude of the effect varied somewhat among groups. We found no statistical effects on distance moved, except for a linetype effect during minute 5 ( $p = 0.0494$ ; HR  $>$  C by  $\sim 12\%$ ) and an exercise  $\times$  diet interaction for minute 3 ( $p = 0.0412$ ). We found no statistical effects on open-field defecation or defecation plus urination (Supplemental Table 1).

### 3.8. Blood lipid profiles

Using bleed delay time and time of day as covariates, adult levels of triglyceride and total cholesterol were unaffected by early-life treatments. Fasted blood glucose levels were increased by early-life exercise (Supplemental Table 1;  $p = 0.0140$ ).

### 3.9. Circulating levels of leptin, adiponectin, and corticosterone

HR mice had approximately 17% greater corticosterone concentration than C mice (Supplemental Table 1;  $p = 0.0068$ ). Among mice exposed to an early-life SD, early-life exercise decreased corticosterone concentration for C mice but increased it for HR

(Supplemental Table 1; exercise  $\times$  diet  $\times$  linetype = 0.0259). Using fat mass as a covariate, early-life exercise treatment increased adult leptin concentration (Supplemental Table 1;  $p = 0.0173$ ), while adiponectin concentration was not affected by linetype or either early-life treatment.

### 3.10. Organ masses

Early-life exercise increased adult triceps surae, subdermal fat pad, and brain mass, but decreased heart ventricle mass (Fig. 8; all  $p \leq 0.0170$ ). Reproductive fat pad and spleen masses were unaffected by either early-life treatment or genetic background. The hearts and brains of HR mice were heavier than those of C mice (all  $p \leq 0.0270$ ).

## 4. Discussion

Physical activity should be a cornerstone of effective prevention and treatment strategies for obesity and its comorbidities (Ryan and Diabetes Prevention Program Research Group 2003; Hill and Wyatt 2005; Janiszewski and Ross 2007, 2009; Church and Blair 2009; Swift et al. 2014; Kirwan et al. 2017). Given the risk of overweight children developing into unhealthy, overweight adults (Malina 2001; Morrison et al. 2007; Singh et al. 2008; The et al. 2010; Lloyd et al. 2012), it is important to attempt to identify early-life factors that might have lasting, positive effects on adult physical activity (Bahls et al. 2014). Implementation of such early-life factors could be much more efficient than waiting until metabolic, cardiovascular, and other disorders appear years later (Vickers et al. 2003; Warden et al. 2007).

The present study is one of the few to examine the possibility of lingering early-life effects of diet and/or physical activity after a substantial washout period with no additional treatment(s). In a full factorial experiment using mice, we tested whether Western diet and/or access to wheels during the juvenile period would: (1) affect juvenile physical activity, caloric intake or body composition; (2) have long-lasting effects into adulthood, following a substantial washout period; and (3) have interactive effects with each other and/or with genetic background. To our knowledge, no other experimental studies have integrated all three factors. We found a number of acute effects of diet and exercise manipulation in juvenile male mice as well as many longer-lasting effects in adults. In

1  
2  
3  
4 addition, some early-life effects interacted with genetic linetype (selectively bred High  
5 Runner vs. non-selected Control lines of mice). Our results have implications for  
6 understanding pre-pubertal environmental effects in humans, including the complex  
7 interactions that result in human obesity and its negative health consequences.  
8  
9

#### 10 11 12 *4.1 Immediate and long-lasting effects of early-life Western diet and exercise* 13 14

##### 15 16 17 4.1.1 Wheel running 18

19 Western diet (WD) increased voluntary exercise (total distance run) during weeks  
20 4 and 5 of the initial period of juvenile exposure (Fig. 2) and also when adults were  
21 retested (Fig. 3), mainly through increased running duration. This effect was greater for  
22 HR mice, and in a previous study was observed only in HR mice (Meek et al. 2010).  
23 Together, these results suggest that inherently athletic individuals and/or those who are  
24 more physically active might gain a greater benefit from high-energy diets.  
25  
26

27 Juvenile exercise increased adult running distance for C mice only (Fig. 3:  
28 exercise  $\times$  linetype  $p = 0.0512$ ). In contrast, Acosta et al. (2015) found that early-life  
29 wheel access increased distance run in both HR and C mice. This difference might relate  
30 to previously an observed seasonal variation in wheel running (see Fig. S4 in Careau et  
31 al. 2013), as the mice in Acosta et al. (2015) were weaned in June whereas ours were  
32 weaned in October. In both studies, early-life wheel effects on adult running distance  
33 were transient in that they became statistically nonsignificant after the first week of adult  
34 testing (Supplemental Table 1).  
35  
36

37 Although the positive effects of juvenile Western diet and exercise that we  
38 observed lasted for only one week, this translates to ~9 months for humans (Demetrius  
39 2006; Flurkey et al. 2007). As these sorts of early-life environmental factors can be  
40 manipulated relatively easily (e.g., through school cafeterias or youth sports programs),  
41 they may be an efficacious tool in the fight against the obesity epidemic and adult  
42 sedentary behavior, although they likely need to be combined with reinforcers during  
43 adulthood (Dalle Grave et al. 2011; Gardner et al. 2012; Strohacker et al. 2014).  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

#### 4.1.2 Body mass and composition

During the juvenile period, male mice given access to wheels did not show the expected reduction in body mass, compared to juveniles without wheel access (e.g. for male mice see Hayes and Williams 1996; Swallow et al. 1999; Acosta et al. 2015). In fact, after three weeks of early-life exposure, mice with wheels had *increased* body mass compared to those without wheels ( $p = 0.0415$ ); this difference was related to increased absolute lean mass ( $p = 0.0104$ ) and accompanied by slightly reduced body fat ( $p = 0.0936$  or  $p = 0.0552$  with lean mass as a covariate).

At the start of adult wheel testing, early-life wheel access had no significant effect on body mass (consistent with Acosta et al. 2015), lean mass or fat mass (see Fig. 4 and Supplemental Table 1), although HR mice weighed significantly less and had lower lean mass as compared with C mice. To our knowledge, no other study has reported effects of early-life exercise on adult body mass after a substantial washout period.

All treatment groups lost body mass across the two weeks of adult wheel testing, but the magnitude of these reductions differed among groups (exercise  $\times$  diet  $\times$  linetype interaction  $p = 0.0059$ ). With respect to body composition, none of the eight experimental groups lost a statistically significant amount of lean mass, whereas all of them lost significant fat mass (see Supplemental Table 1). Consistent with a previous study that used a 6-day wheel test, HR lines lost less fat mass than C mice during adult wheel testing (see Supplemental Table 1) (Hiramatsu and Garland 2018), indicating that the regulation of body weight when faced with acute exercise varies with genetic background.

Although WD consistently increased fat mass during the three weeks of early-life exposure, simultaneous exercise partially protected mice from this increase (though it was not eliminated completely [(exercise  $\times$  diet  $p = 0.0170$  for Week 1,  $p = 0.5136$  for Week 2, and  $p = 0.0355$  for Week 3)]). This effect mostly faded by the end of the 8-week washout, during which all mice received SD. Early-life WD increased adult lean mass ( $p = 0.0316$ ), but the effect disappeared after two weeks of adult exercise ( $p = 0.0960$ ).

#### 4.1.3 Caloric intake

We expected that body mass-adjusted caloric intake would increase with wheel access (Swallow et al. 2001; Acosta et al. 2015; Copes et al. 2015). Surprisingly, in the first week of juvenile treatments, early-life wheel access decreased caloric intake for mice

feeding on WD and increased it for mice feeding on SD (Fig. 5). During this first week, mice are experiencing multiple environmental changes (novel exposures), including not only diet and exercise opportunity, but also single housing, which affects social behavior and thermoregulation, so their normal homeostatic mechanisms (both physiological and neurobiological) may not yet have adjusted. During the second and third weeks, however, wheel access had the expected positive effect on caloric intake and did not interact with diet or linetype. In any case, the neural and endocrine mechanisms that underlie homeostatic responses to novel (acute) versus longer-term (chronic) environmental changes deserve further study.

During the first week of adult wheel testing, HR mice had greater mass-adjusted caloric intake than C mice, consistent with previous studies (e.g., Acosta et al. 2015; Copes et al. 2015; Hiramatsu and Garland 2018), with no statistical effects of either early-life WD ( $p = 0.0929$ ) or exercise (Fig. 6). During the second week of adult testing, an effect of early-life exercise emerged, in which caloric intake was decreased in HR mice, but not in C mice (exercise  $\times$  linetype  $p = 0.0126$ ). When we added distance run as a covariate, to account for the greater running by HR mice, the interaction disappeared, and the effect of early-life exercise was negative for both HR and C mice (Supplemental Table 1). Given that early-life exercise increased adult wheel running in both HR and C mice, and that HR mice ran much more than C, this effect suggests an alteration in digestive or metabolic efficiency, or perhaps in the regulation of body weight, although the loss in body mass across the two weeks of adult wheel access did not show any statistically significant interactive effects (Supplemental Table 1). In any case, early-life exercise exposure combined with adult exercise had a dampening effect on caloric intake, which could have important implications for human health (Chin et al. 2016; Golbidi et al. 2017; Obert et al. 2017).

#### 4.1.4 Western diet and sucrose preference

As adults, almost all mice preferred the WD to standard diet when given 24-hour access to both diets, as is commonly reported (e.g., Acosta et al. 2017). On average, however, those given WD as juveniles had a significantly stronger preference for WD than those given juvenile SD (Supplemental Table 1). Although WD is high in sugar and fat, we did not find any statistically significant effects on adult sucrose preference,

1  
2  
3  
4 suggesting that familiarity with the taste of the fatty component of WD may be driving the  
5 diet preference. Additionally, a three-way interaction effect on WD preference was found  
6 (exercise  $\times$  diet  $\times$  linetype  $p = 0.0578$ ), indicating some effect of genetic background.  
7 Although few rodent studies are available (e.g., Teegarden et al. 2009), various human  
8 studies indicate that early-life factors can affect adult dietary preferences (e.g., Lussana  
9 et al. 2008; Paglia 2019).  
10  
11  
12  
13  
14

#### 15 16 4.1.5 Open-field behavior 17

18 Early-life wheel access generally reduced adult anxiety-like behavior (i.e., less  
19 time spent near walls in the open field), although the strength of the effect varied among  
20 groups (exercise  $\times$  diet  $\times$  linetype  $p = 0.0432$ ). In principle, effects on anxiety-like  
21 behavior might be mediated by plasma corticosterone levels (Korte 2001; McEwen et al.  
22 2015). As noted in the next section, adult plasma corticosterone levels also showed a 3-  
23 way interaction (exercise  $\times$  diet  $\times$  linetype  $p = 0.0259$ ). The patterns for these 3-way  
24 interactions were somewhat different, but the correlation of the eight group means for  
25 time spent near walls (over all 5 minutes, squared) and plasma corticosterone (log-  
26 transformed) was positive (Pearson's  $r = 0.546$ ,  $p = 0.1615$ ), suggesting that  
27 corticosterone might have played some role in mediating effects of early-life wheel  
28 access on anxiety-like behavior. The reduction in anxiety-like behaviors cannot be  
29 attributed to effects of early-life wheel access on activity levels because distance moved  
30 in the open-field test was not affected in a way that paralleled time spent near walls. An  
31 interesting direction for future studies would be to evaluate a possible role of the  
32 neuropeptide corticotropin-releasing hormone, which regulates corticosterone secretion,  
33 strongly influences anxiety-like behavior, and has been suggested to mediate effects of  
34 experience (stress) during the juvenile period on adult phenotypes (Burke and Miczek  
35 2014; Syed et al. 2017; Zhang et al. 2017).  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52

#### 53 4.1.6 Leptin, adiponectin, corticosterone, and blood glucose 54

55 Leptin is produced by white adipose tissue and, among other functions, regulates  
56 body weight by increasing energy expenditure and inhibiting food intake (Zhang et al.  
57 2005; Triantafyllou et al. 2016; Zhang and Chua 2017), along with a host of other effects,  
58 including an important role in glucose metabolism (Meek and Morton 2016). As expected  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 from numerous previous studies, including those involving the HR mice (Girard et al.  
5 2007; Acosta et al. 2015; Hiramatsu et al. 2017), body fat was a strong positive predictor  
6 of adult levels of circulating leptin in the present study. In addition, early-life wheel  
7 access increased fat-adjusted leptin levels in both linetypes, whereas Acosta et al (2015)  
8 found that early-life wheel access interacted with linetype (no wheels: HR>C; wheels:  
9 HR<C). The discrepancy in these results may be explainable by Acosta et al.'s. use of  
10 visceral fat pad mass and our use of body fat measured by MRI as covariates: dissected  
11 visceral fat pads are only a subset of a mouse's leptin-producing adipocytes; however,  
12 not all body fat measured via MRI produces leptin (Cinti 2012). It is also worth noting that  
13 leptin regulation in mice can be influenced by age (Ahren et al. 1997); we collected  
14 terminal serum samples when mice were 20 weeks old, whereas Acosta et al. (2015)  
15 took samples at ~15 weeks.

16  
17 Although some previous studies have reported higher plasma adiponectin levels in  
18 HR mice as compared with C lines (Vanholt et al. 2007, 2008; Garland, Jr. et al. 2017),  
19 we did not find any statistically significant effects on adiponectin. However, adult plasma  
20 corticosterone showed a 3-way interaction (exercise x diet x linetype  $p = 0.0259$ ), as well  
21 as a main effect of linetype ( $p = 0.0068$ ). The selected HR lines always have elevated  
22 corticosterone levels, as has generally been found in previous studies (e.g., Malisch et al.  
23 2007, 2008, 2016; Downs et al. 2012; but see Vanholt et al. 2007; Hiramatsu et al.  
24 2017). In addition, HR mice exposed to early-life wheel access and SD had much higher  
25 levels of corticosterone than any other group (Supplemental Table 1).

26  
27 To our knowledge, no previous studies have tested for long-lasting effects of early-  
28 life exercise on adult blood glucose concentrations. We observed higher glucose levels  
29 in mice given juvenile wheel access than in those housed without wheels.  
30 Perhaps this effect is related to the effects observed when humans and rodents are  
31 cycled between standard and high-fat diets (e.g., Strohacker et al. 2009). It is also  
32 possible that early-life exercise has affected glucose tolerance (Oelkrug et al. 2020);  
33 however, the current study did not include a glucose tolerance test, which may have been  
34 useful in addressing these questions. In any case, the early-life exercise effects on leptin  
35 noted above may be involved in the effects on blood glucose (Meek and Morton 2016).



#### 4.1.7 Organ masses

Numerous studies spanning many decades show that providing laboratory rodents with chronic wheel access leads to changes in the size of internal organs relative to body size, but few have tested for long-lasting effects of juvenile exercise alone. In a previous study of male HR and C mice, Acosta et al. (2015) found no effects of early-life wheel access on visceral fat pads, heart ventricles, liver, or spleen masses. Some of our results are similar, but others differ. Heart ventricle and liver masses were affected by multiple interactions among diet, exercise, and linetype (see Fig. 8A, C). These effects could be either beneficial or detrimental, depending on the cause. For example, an increase in liver mass may be detrimental if caused primarily by WD (Fig. 8D), which often causes cirrhosis or non-alcoholic fatty liver disease (Zhou et al. 2014; Maurice and Manousou 2018, respectively).

An apparently novel finding was the positive effect of early-life exercise on calf-muscle mass for all groups (Fig. 8B). This type of effect, if it occurs in humans, has obvious relevance for human health from early-life to advanced age (Westcott 2012; McLeod et al. 2016; Orsso et al. 2019).

Another notable effect of early-life exercise was an increased brain mass in all groups. Several recent studies have shown that chronic exercise can increase brain mass in rodents, even within one week (Sumiyoshi et al. 2014). To our knowledge, however, this is the first report of a juvenile effect on brain mass that persists into adulthood after a substantial washout period. Exercise is known to upregulate growth factors, including brain-derived neurotrophic factor, which may have an even larger effect during a developmental period (e.g., see Park and Höke 2014 for rodents and Negaresh et al. 2018 for humans).

#### 4.2 Differences between High Runner and non-selected Control lines

The current study reaffirms many previously reported differences between the HR and C linetypes (Acosta et al. 2015; Garland, Jr. et al. 2016; Wallace and Garland, Jr. 2016; Hiramatsu et al. 2017). Average daily wheel-running distance during juvenile testing was higher for HR than C mice, due mainly to greater average running speeds, consistent with many previous studies (e.g., Swallow et al. 1999; Meek et al. 2010; Acosta et al. 2015) (Fig. 2A).



1  
2  
3  
4 In addition, as adults, HR mice were smaller in total body mass and had leaner  
5 bodies, and (with body mass as a covariate), they had greater maximal aerobic capacity,  
6 heavier hearts and brains, and elevated circulating levels of corticosterone. Other  
7 differences, including several that involved interactions with early-life diet and/or exercise  
8 exposure, are mentioned above.  
9  
10  
11  
12  
13

#### 14 4.3 Characteristics of mini-muscle individuals

15 As described above (see 2.3.9. *Statistical analysis*), the mini-muscle phenotype  
16 and underlying gene were an unexpected discovery in the High Runner mouse selection  
17 experiment (Garland, Jr. et al. 2002; Kelly et al. 2013). In addition to their greatly  
18 reduced hindlimb muscle mass, the present study confirmed several previously reported  
19 differences between mini- and normal-muscle individuals, including heavier heart  
20 ventricles, livers, and greater VO<sub>2</sub>max (Rezende 2006; Kolb et al. 2010; Hiramatsu et al.  
21 2017; Kelly et al. 2017; Kay et al. 2019). In addition, we found some new differences,  
22 including that mini-muscle individuals spent more time near the walls in an open-field  
23 behavior test (suggesting elevated anxiety) and increased total body fat. Future studies  
24 will be required to determine the molecular and cellular pathways through which the mini-  
25 muscle gene causes such a wide range of pleiotropic effects.  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39

## 40 5. Limitations and Concluding Remarks

41  
42  
43 One limitation of the present study is that all mice were given two weeks of adult  
44 wheel access, followed by one day with the wheels removed, prior to dissection.  
45 Therefore, organ masses and circulating concentrations of adiponectin and leptin would  
46 have been affected by early-life effects, two weeks of training effects (physical  
47 conditioning), plus any acute "rebound" effects that might have occurred over ~24 hours  
48 (e.g., see Dumke et al. 2001). Any of these effects could have obscured the others. In  
49 future experiments, we intend to have multiple cohorts of subjects, such as one that is  
50 sampled prior to adult wheel testing.  
51  
52  
53  
54  
55  
56  
57

58 Another limitation is that we studied only males. Importantly, nutrient intake in  
59 early life has been shown to have sex-specific effects on the regulation of physical  
60  
61  
62  
63  
64  
65

activity (i.e., wheel running) in mice (Leszczynski et al. 2019). In an early-life experiment on the same mouse model as the current study, Hiramatsu et al. (2017) observed sex differences as well (see their Fig. 8 and online supplemental Fig. 4). Moreover, given that many aspects of physical activity (including both motivation and ability), exercise physiology, body composition, and hormone profiles are sexually dimorphic in both mice (Wells 2007; Lightfoot 2008; Palmer and Clegg 2015; Klinker et al. 2017; van Keulen et al. 2020) and humans (Buffa et al. 2001; Malisch et al. 2007; Ingvorsen et al. 2017; Rosenfeld 2017; Zore et al. 2018), future studies should include both sexes.

Considering only statistically significant main effects ( $p < 0.05$ ), we observed 11 effects of early-life exercise but only three effects of early-life Western diet on adult traits, which suggests that the former may be a more effective target for interventions with human youth. Evidence concerning whether activity levels of children can be increased in particular areas without compensation in others is the subject of ongoing debate over a proposed homeostatic compensatory mechanism known as an “activity-stat” (Rowland 1998; Wilkin 2011; Gomersall et al. 2016). In any case, studies of declining physical activity in children suggest that sedentary behaviors begin displacing physical activity habits much earlier than is generally believed and that interventions should begin prior to initial school admission (Reilly 2011, 2016).

Our results may also be relevant to the current COVID-19 pandemic, wherein the lack of regular physical activity caused by lockdowns and other restrictions on mobility (and likely effects on diet) is becoming a widespread epidemiological issue (Mattioli et al. 2020). In particular, children are poised to develop simultaneous lifestyle habits of physical inactivity and high consumption of calorie-dense foods in quarantine that may have lasting effects into adulthood, such as overweight and obesity (Rundle et al. 2020). (Of course, SARS-CoV-2 infection itself may be viewed as an early-life environmental factor with potentially long-lasting adverse effects.) Changes in lifestyle habits related to diet and exercise appear to differ by country. For example, Di Renzo et al. (2020) found increased physical exercise and consumption of Mediterranean diet in Italy during the pandemic, whereas Sidor and Rzymiski (2020) found increased consumption of fast foods (which are usually high-fat, high-sugar) in Poland. In a survey of adults in Australia, 48.9% of respondents reported reduced daily physical activity as a result of pandemic confinement (Stanton et al. 2020). In another survey conducted on young people (aged

10 to 19 years) in a number of South American and European countries, respondents reported a significant increase in the intake of foods high in fat and sugar since the pandemic began (Ruiz-Roso et al. 2020). Because the pandemic is fairly recent, long-term effects of these lifestyle changes are yet unknown; however, we predict that the COVID-19 pandemic will have long-lasting effects on physical activity and food preferences as today's affected children become adults. Given the acute maladies associated with physical inactivity and the virus itself (Dunton et al. 2020; Woods et al. 2020), it seems especially important to implement interventions that may engender better lifestyle habits in children, who now experience the dietary and physical activity challenges that come with school closures and reduced access to recreational activities (Rundle et al. 2020; Wang et al. 2020).

## Conflict of interest statement

The authors declare no conflict of interest.

## Acknowledgments

This work was supported US NIH grant R21HD084856 to T.G. and Wendy Saltzman, in part by US NSF grant DEB-1655362 to T.G., and in part by US USDA project CA-R-EEOB-5205-H to T.G. We thank M. A. Chappell for technical and methodological assistance, and T. H. Meek helpful discussions.

## 6. References

- Acosta, W., T. H. Meek, H. Schutz, E. M. Dlugosz, and T. Garland, Jr. 2017. Preference for Western diet coadapts in High Runner mice and affects voluntary exercise and spontaneous physical activity in a genotype-dependent manner. *Behav. Processes* 135:56–65.
- Acosta, W., T. H. Meek, H. Schutz, E. M. Dlugosz, K. T. Vu, and T. Garland. 2015. Effects of early-onset voluntary exercise on adult physical activity and associated phenotypes in mice. *Physiol. Behav.* 149:279–286.
- Adam, T. C., and E. S. Epel. 2007. Stress, eating and the reward system. *Physiol. Behav.* 91:449–458.
- Ahren, B., S. Mansson, R. L. Gingerich, and P. J. Havel. 1997. Regulation of plasma leptin in mice: influence of age, high-fat diet, and fasting. *Am. J. Physiol.-Regul. Integr. Comp. Physiol.* 273:R113–R120. American Physiological Society.
- Andersen, L. G., L. Ångquist, M. Gamborg, L. Byberg, C. Bengtsson, D. Canoy, J. G. Eriksson, M. Eriksson, M.-R. Järvelin, L. Lissner, T. I. Nilsen, M. Osler, K. Overvad, F. Rasmussen, M. K. Salonen, L. Schack-Nielsen, T. H. Tammelin, T.-P. Tuomainen, T. I. A. Sørensen, J. L. Baker, and for the N. S. Group. 2009. Birth weight in relation to leisure time physical activity in adolescence and adulthood: meta-analysis of results from 13 nordic cohorts. *PLOS ONE* 4:e8192. Public Library of Science.
- Avenell, A., T. J. Brown, M. A. McGee, M. K. Campbell, A. M. Grant, J. Broom, R. T. Jung, and W. C. S. Smith. 2004. What interventions should we add to weight reducing diets in adults with obesity? A systematic review of randomized

controlled trials of adding drug therapy, exercise, behaviour therapy or combinations of these interventions. *J. Hum. Nutr. Diet. Off. J. Br. Diet. Assoc.* 17:293–316.

Bahls, M., R. D. Sheldon, P. Taheripour, K. A. Clifford, K. B. Foust, E. D. Breslin, J. N. Marchant-Forde, R. A. Cabot, M. Harold Laughlin, C. A. Bidwell, and S. C. Newcomer. 2014. Mother's exercise during pregnancy programmes vasomotor function in adult offspring. *Exp. Physiol.* 99:205–219.

Baker, J., M. Workman, E. Bedrick, M. A. Frey, M. Hurtado, and O. Pearson. 2010. Brains versus brawn: An empirical test of Barker's brain sparing model. *Am. J. Hum. Biol.* 22:206–215.

Baker, M. S., G. Li, J. J. Kohorst, and R. A. Waterland. 2015. Fetal growth restriction promotes physical inactivity and obesity in female mice. *Int. J. Obes.* 2005 39:98–104.

Barker, D. J. P. 2003. The developmental origins of adult disease. *Eur. J. Epidemiol.* 18:733–736.

Blair, S. N., and J. N. Morris. 2009. Healthy hearts--and the universal benefits of being physically active: physical activity and health. *Ann. Epidemiol.* 19:253–256.

Blondin, S. A., S. Anzman-Frasca, H. C. Djang, and C. D. Economos. 2016. Breakfast consumption and adiposity among children and adolescents: an updated review of the literature. *Pediatr. Obes.* 11:333–348.

Boitard, C., N. Etchamendy, J. Sauvant, A. Aubert, S. Tronel, A. Marighetto, S. Layé, and G. Ferreira. 2012. Juvenile, but not adult exposure to high-fat diet impairs relational memory and hippocampal neurogenesis in mice. *Hippocampus* 22:2095–2100.

- 1  
2  
3 Bouchard, C., T. Rankinen, and J. A. Timmons. 2011. Genomics and genetics in the  
4  
5 biology of adaptation to exercise. *Compr. Physiol.* 1:1603–1648.  
6  
7  
8 Breier, B. H., S. O. Krechowec, and M. H. Vickers. 2006. Programming of obesity—  
9  
10 experimental evidence. Pp. 145–156 *in* E. M. Wintour and J. A. Owens, eds. *Early*  
11  
12 *life origins of health and disease*. Springer US, Boston, MA.  
13  
14  
15 Brennehan, D. E., and C. O. Rutledge. 1982. Effect of dietary lipid on locomotor activity  
16  
17 and response to psychomotor stimulants. *Psychopharmacology (Berl.)* 76:260–  
18  
19 264.  
20  
21  
22 Britton, S. L., and L. G. Koch. 2001. Animal genetic models for complex traits of physical  
23  
24 capacity. *Exerc. Sport Sci. Rev.* 29:7–14.  
25  
26  
27 Bronikowski, A. M., P. A. Carter, J. G. Swallow, I. A. Girard, J. S. Rhodes, and T.  
28  
29 Garland. 2001. Open-field behavior of house mice selectively bred for high  
30  
31 voluntary wheel-running. *Behav. Genet.* 31:309–316.  
32  
33  
34 Bronikowski, A. M., T. J. Morgan, T. Garland, Jr., and P. A. Carter. 2006. The evolution of  
35  
36 aging and age-related physical decline in mice selectively bred for high voluntary  
37  
38 exercise. *Evolution* 60:1494–1508.  
39  
40  
41  
42 Buffa, R., E. Marini, and G. Floris. 2001. Variation in sexual dimorphism in relation to  
43  
44 physical activity. *Am. J. Hum. Biol. Off. J. Hum. Biol. Counc.* 13:341–348.  
45  
46  
47 Burke, A. R., and K. A. Miczek. 2014. Stress in adolescence and drugs of abuse in rodent  
48  
49 models: Role of dopamine, CRF, and HPA axis. *Psychopharmacology (Berl.)*  
50  
51 231:1557–1580.  
52  
53  
54 Careau, V., M. E. Wolak, P. A. Carter, and T. Garland. 2013. Limits to behavioral  
55  
56 evolution: the quantitative genetics of a complex trait under directional selection:  
57  
58 quantitative genetics of a selection limit. *Evolution* 67:3102–3119.  
59  
60  
61  
62  
63  
64  
65

- Chin, S.-H., C. N. Kahathuduwa, and M. Binks. 2016. Physical activity and obesity: what we know and what we need to know. *Obes. Rev. Off. J. Int. Assoc. Study Obes.* 17:1226–1244.
- Choh, A. C., E. W. Demerath, M. Lee, K. D. Williams, B. Towne, R. M. Siervogel, S. A. Cole, and S. A. Czerwinski. 2009. Genetic analysis of self-reported physical activity and adiposity: the southwest ohio family study. *Public Health Nutr.* 12:1052–1060.
- Church, T. S., and S. N. Blair. 2009. When will we treat physical activity as a legitimate medical therapy...even though it does not come in a pill? *Br. J. Sports Med.* 43:80–81.
- Cinti, S. 2012. The adipose organ at a glance. *Dis. Model. Mech.* 5:588–594.
- Claghorn, G. C., Z. Thompson, K. Wi, L. Van, and T. Garland. 2017. Caffeine stimulates voluntary wheel running in mice without increasing aerobic capacity. *Physiol. Behav.* 170:133–140.
- Coccurello, R., F. R. D'Amato, and A. Moles. 2009. Chronic social stress, hedonism and vulnerability to obesity: lessons from rodents. *Neurosci. Biobehav. Rev.* 33:537–550.
- Copes, L. E., H. Schutz, E. M. Dlugosz, W. Acosta, M. A. Chappell, and T. Garland. 2015. Effects of voluntary exercise on spontaneous physical activity and food consumption in mice: Results from an artificial selection experiment. *Physiol. Behav.* 149:86–94.
- Cornier, M.-A., D. Dabelea, T. L. Hernandez, R. C. Lindstrom, A. J. Steig, N. R. Stob, R. E. Van Pelt, H. Wang, and R. H. Eckel. 2008. The metabolic syndrome. *Endocr. Rev.* 29:777–822.

- 1  
2  
3  
4 Dai, Y., S. Thamocharan, M. Garg, B.-C. Shin, and S. U. Devaskar. 2012.  
5  
6 Superimposition of postnatal calorie restriction protects the aging male intrauterine  
7  
8 growth- restricted offspring from metabolic maladaptations. *Endocrinology*  
9  
10 153:4216–4226.  
11  
12  
13 Dalle Grave, R., S. Calugi, E. Centis, M. El Ghoch, and G. Marchesini. 2011. Cognitive-  
14  
15 behavioral strategies to increase the adherence to exercise in the management of  
16  
17 obesity. *J. Obes.* 2011:1–11.  
18  
19  
20 de Boo, H. A., and J. E. Harding. 2006. The developmental origins of adult disease  
21  
22 (Barker) hypothesis. *Aust. N. Z. J. Obstet. Gynaecol.* 46:4–14.  
23  
24  
25 Demetrius, L. 2006. Aging in mouse and human systems: a comparative study. *Ann. N.*  
26  
27 *Y. Acad. Sci.* 1067:66–82.  
28  
29  
30 Desai, M., T. Li, G. Han, and M. G. Ross. 2014. Programmed hyperphagia secondary to  
31  
32 increased hypothalamic SIRT1. *Brain Res.* 1589:26–36.  
33  
34  
35 Di Renzo, L., P. Gualtieri, F. Pivari, L. Soldati, A. Attinà, G. Cinelli, C. Leggeri, G.  
36  
37 Caparello, L. Barrea, F. Scerbo, E. Esposito, and A. De Lorenzo. 2020. Eating  
38  
39 habits and lifestyle changes during COVID-19 lockdown: an Italian survey. *J.*  
40  
41 *Transl. Med.* 18.  
42  
43  
44 Dishman, R. K., J. F. Sallis, and D. R. Orenstein. 1985. The determinants of physical  
45  
46 activity and exercise. *Public Health Rep.* 100:158–171.  
47  
48  
49 Dlugosz, E. M., M. A. Chappell, D. G. McGillivray, D. A. Syme, and T. Garland. 2009.  
50  
51 Locomotor trade-offs in mice selectively bred for high voluntary wheel running. *J.*  
52  
53 *Exp. Biol.* 212:2612–2618.  
54  
55  
56 Dohle, S., and B. Wansink. 2013. Fit in 50 years: participation in high school sports best  
57  
58 predicts one's physical activity after Age 70. *BMC Public Health* 13:1100.  
59  
60  
61  
62  
63  
64  
65



- 1  
2  
3  
4 Donovan, E., C. E. Hernandez, L. R. Matthews, M. Oliver, A. Jaquiere, F. Bloomfield, and  
5  
6 J. Harding. 2013. Periconceptional undernutrition in sheep leads to decreased  
7  
8 locomotor activity in a natural environment. *J. Dev. Orig. Dis.*  
9
- 10  
11 Downing, C., T. E. Johnson, C. Larson, T. I. Leakey, R. N. Siegfried, T. M. Rafferty, and  
12  
13 C. A. Cooney. 2011. Subtle decreases in dna methylation and gene expression at  
14  
15 the mouse *igf2* locus following prenatal alcohol exposure: effects of a methyl-  
16  
17 supplemented diet. *Alcohol Fayettev. N* 45:65–71.  
18  
19
- 20  
21 Downs, C. J., H. Schutz, T. H. Meek, E. M. Dlugosz, W. Acosta, K. S. de Wolski, J. L.  
22  
23 Malisch, J. P. Hayes, and T. Garland, Jr. 2012. Within-lifetime trade-offs but  
24  
25 evolutionary freedom for hormonal and immunological traits: evidence from mice  
26  
27 bred for high voluntary exercise. *J. Exp. Biol.* 215:1651–1661.  
28  
29
- 30  
31 Drenowatz, C., J. J. Carlson, K. A. Pfeiffer, and J. C. Eisenmann. 2012. Joint association  
32  
33 of physical activity/screen time and diet on CVD risk factors in 10-year-old  
34  
35 children. *Front. Med.* 6:428–435.  
36  
37
- 38  
39 Dumke, C. L., J. S. Rhodes, T. Garland, Jr., E. Maslowski, J. G. Swallow, A. C. Wetter,  
40  
41 and G. D. Cartee. 2001. Genetic selection of mice for high voluntary wheel  
42  
43 running: effect on skeletal muscle glucose uptake. *J. Appl. Physiol.* 91:1289–1297.  
44
- 45  
46 Dunton, G. F., B. Do, and S. D. Wang. 2020. Early effects of the COVID-19 pandemic on  
47  
48 physical activity and sedentary behavior in children living in the U.S. *BMC Public*  
49  
50 *Health* 20:1351.  
51
- 52  
53 Eisenmann, J. C. 2006. Insight into the causes of the recent secular trend in pediatric  
54  
55 obesity: Common sense does not always prevail for complex, multi-factorial  
56  
57 phenotypes. *Prev. Med.* 42:329–335.  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2  
3 Eisenmann, J. C. 2004. Physical activity and cardiovascular disease risk factors in  
4  
5 children and adolescents: an overview. *Can. J. Cardiol.* 20:295–301.  
6  
7  
8 Esposito, K., M. I. Maiorino, A. Ceriello, and D. Giugliano. 2010. Prevention and control of  
9  
10 type 2 diabetes by Mediterranean diet: a systematic review. *Diabetes Res. Clin.*  
11  
12 *Pract.* 89:97–102.  
13  
14  
15 Flegal, K. M., C. L. Ogden, J. A. Yanovski, D. S. Freedman, J. A. Shepherd, B. I.  
16  
17 Graubard, and L. G. Borrud. 2010. High adiposity and high body mass index-for-  
18  
19 age in US children and adolescents overall and by race-ethnic group. *Am. J. Clin.*  
20  
21 *Nutr.* 91:1020–1026.  
22  
23  
24  
25 Flurkey, K., J. Curren, and D. Harrison. 2007. Mouse models in aging research. *Fac. Res.*  
26  
27 637–672.  
28  
29  
30 Frazier, C. R. M., P. Mason, X. Zhuang, and J. A. Beeler. 2008. Sucrose exposure in  
31  
32 early life alters adult motivation and weight Gain. *PLOS ONE* 3:e3221.  
33  
34  
35 Fuss, J., N. M.-B. Ben Abdallah, M. A. Vogt, C. Touma, P. G. Pacifici, R. Palme, V.  
36  
37 Witzemann, R. Hellweg, and P. Gass. 2009. Voluntary exercise induces anxiety-  
38  
39 like behavior in adult C57BL/6J mice correlating with hippocampal neurogenesis.  
40  
41 *Hippocampus* 364–376.  
42  
43  
44  
45 Gardner, B., P. Lally, and J. Wardle. 2012. Making health habitual: the psychology of  
46  
47 ‘habit-formation’ and general practice. *Br. J. Gen. Pract.* 62:664–666.  
48  
49  
50 Gardner, D. S., and P. Rhodes. 2009. Developmental origins of obesity: programming of  
51  
52 food intake or physical activity? *Adv. Exp. Med. Biol.* 646:83–93.  
53  
54  
55 Garland, Jr., T., M. D. Cadney, and R. A. Waterland. 2017. Early-life effects on adult  
56  
57 physical activity: concepts, relevance, and experimental approaches. *Physiol.*  
58  
59 *Biochem. Zool.* 90:1–14.  
60  
61  
62  
63  
64  
65

- Garland, Jr., T., M. T. Morgan, J. G. Swallow, J. S. Rhodes, I. Girard, J. G. Belter, and P. A. Carter. 2002. Evolution of a small-muscle polymorphism in lines of house mice selected for high activity levels. *Evolution* 56:1267–1275.
- Garland, Jr., T., and M. R. Rose (eds). 2009. *Experimental Evolution: Concepts, Methods, and Applications of Selection Experiments*. First Edition. University of California Press, Berkeley.
- Garland, Jr., T., H. Schutz, M. A. Chappell, B. K. Keeney, T. H. Meek, L. E. Copes, W. Acosta, C. Drenowatz, R. C. Maciel, G. van Dijk, C. M. Kotz, and J. C. Eisenmann. 2011. The biological control of voluntary exercise, spontaneous physical activity and daily energy expenditure in relation to obesity: human and rodent perspectives. *J. Exp. Biol.* 214:206–229.
- Garland, Jr., T., M. Zhao, and W. Saltzman. 2016. Hormones and the Evolution of Complex Traits: Insights from Artificial Selection on Behavior. *Integr. Comp. Biol.* 56:207–224.
- Girard, I., E. L. Rezende, and T. Garland Jr. 2007. Leptin levels and body composition of mice selectively bred for high voluntary locomotor activity. *Physiol. Biochem. Zool.* 80:568–579.
- Gluckman, P. D., M. A. Hanson, and A. S. Beedle. 2007. Early life events and their consequences for later disease: A life history and evolutionary perspective. *Am. J. Hum. Biol.* 19:1–19.
- Golbidi, S., A. Daiber, B. Korac, H. Li, M. F. Essop, and I. Laher. 2017. Health benefits of fasting and caloric restriction. *Curr. Diab. Rep.* 17:123.
- Golde, W. T., P. Gollobin, and L. L. Rodriguez. 2005. A rapid, simple, and humane method for submandibular bleeding of mice using a lancet. *Lab Anim.* 34:39–43.

- Gomersall, S. R., C. Maher, C. English, A. V. Rowlands, J. Dollman, K. Norton, and T. Olds. 2016. Testing the activitystat hypothesis: a randomised controlled trial. *BMC Public Health* 16:900.
- Graff, M., K. E. North, K. L. Monda, E. M. Lange, L. A. Lange, G. Guo, and P. Gordon-Larsen. 2011. The combined influence of genetic factors and sedentary activity on body mass changes from adolescence to young adulthood: the national longitudinal adolescent health study. *Diabetes Metab. Res. Rev.* 27:63–69.
- Guderley, H., P. Houle-Leroy, G. M. Diffie, D. M. Camp, and T. Garland, Jr. 2006. Morphometry, ultrastructure, myosin isoforms, and metabolic capacities of the “mini muscles” favoured by selection for high activity in house mice. *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* 144:271–282.
- Hallal, P. C., J. C. K. Wells, F. F. Reichert, L. Anselmi, and C. G. Victora. 2006. Early determinants of physical activity in adolescence: prospective birth cohort study. *BMJ* 332:1002–1007.
- Halpern, A., M. C. Mancini, M. E. C. Magalhães, M. Fisberg, R. Radominski, M. C. Bertolami, A. Bertolami, M. E. de Melo, M. T. Zanella, M. S. Queiroz, and M. Nery. 2010. Metabolic syndrome, dyslipidemia, hypertension and type 2 diabetes in youth: from diagnosis to treatment. *Diabetol. Metab. Syndr.* 2:55.
- Hannon, R. M., S. A. Kelly, K. M. Middleton, E. M. Kolb, D. Pomp, and T. Garland, Jr. 2008. Phenotypic effects of the “mini-muscle” allele in a large HR x C57BL/6J mouse backcross. *J. Hered.* 99:349–354.
- Hanson, M. A., and P. D. Gluckman. 2008. Developmental origins of health and disease: new Insights. *Basic Clin. Pharmacol. Toxicol.* 102:90–93.

- Harris, K. C., L. K. Kuramoto, M. Schulzer, and J. E. Retallack. 2009. Effect of school-based physical activity interventions on body mass index in children: a meta-analysis. *CMAJ Can. Med. Assoc. J. J. Assoc. Medicales Can.* 180:719–726.
- Hayes, A., and D. A. Williams. 1996. Beneficial effects of voluntary wheel running on the properties of dystrophic mouse muscle. *J. Appl. Physiol.* 80:670–679.
- Hill, J. O., and H. R. Wyatt. 2005. Role of physical activity in preventing and treating obesity. *J. Appl. Physiol.* 99:765–770.
- Hillis, D. A., L. Yadgary, G. M. Weinstock, F. P.-M. de Villena, D. Pomp, A. S. Fowler, S. Xu, F. Chan, and T. Garland, Jr. 2020. Genetic basis of aerobically supported voluntary exercise: results from a selection experiment with house mice. *Genetics* 216:781–804.
- Hills, A. P., L. B. Andersen, and N. M. Byrne. 2011. Physical activity and obesity in children. *Br. J. Sports Med.* 45:866–870.
- Hiramatsu, L., and T. Garland, Jr. 2018. Mice selectively bred for high voluntary wheel-running behavior conserve more fat despite increased exercise. *Physiol. Behav.* 194:1–8.
- Hiramatsu, L., J. C. Kay, Z. Thompson, J. M. Singleton, G. C. Claghorn, R. L. Albuquerque, B. Ho, B. Ho, G. Sanchez, and T. Garland, Jr. 2017. Maternal exposure to Western diet affects adult body composition and voluntary wheel running in a genotype-specific manner in mice. *Physiol. Behav.* 179:235–245.
- Houle-Leroy, P., T. Garland, Jr., J. G. Swallow, and H. Guderley. 2000. Effects of voluntary activity and genetic selection on muscle metabolic capacities in house mice *Mus domesticus*. *J. Appl. Physiol.* 89:1608–1616.

- 1  
2  
3  
4 Ingvorsen, C., N. A. Karp, and C. J. Lelliott. 2017. The role of sex and body weight on the  
5  
6 metabolic effects of high-fat diet in C57BL/6N mice. *Nutr. Diabetes* 7:e261.  
7  
8 Janiszewski, P. M., and R. Ross. 2007. Physical activity in the treatment of obesity:  
9  
10 beyond body weight reduction. *Appl. Physiol. Nutr. Metab. Physiol. Appl. Nutr.*  
11  
12 *Metab.* 32:512–522.  
13  
14 Janiszewski, P. M., and R. Ross. 2009. The utility of physical activity in the management  
15  
16 of global cardiometabolic risk. *Obes. Silver Spring Md* 17 Suppl 3:S3–S14.  
17  
18  
19 Janssen, I., V. Carson, I.-M. Lee, P. T. Katzmarzyk, and S. N. Blair. 2013. Years of life  
20  
21 gained due to leisure-time physical activity in the U.S. *Am. J. Prev. Med.* 44:23–  
22  
23 29.  
24  
25  
26  
27 Jónás, I., L. M. Vaanholt, M. Doornbos, T. Garland, Jr., A. J. W. Scheurink, C. Nyakas,  
28  
29 and G. van Dijk. 2010. Effects of selective breeding for increased wheel-running  
30  
31 behavior on circadian timing of substrate oxidation and ingestive behavior. *Physiol.*  
32  
33 *Behav.* 99:549–554.  
34  
35  
36  
37 Jung, A., T. Curtis, M. Turner, and J. Lightfoot. 2006. Influence of age of exposure to a  
38  
39 running wheel on activity in inbred mice. *Med. Sci. Sports Exerc.* 38:51–56.  
40  
41  
42 Jung, A. P., and D. R. Luthin. 2010. Wheel access does not attenuate weight gain in mice  
43  
44 fed high-fat or high-CHO diets. *Med. Sci. Sports Exerc.* 42:355–360.  
45  
46  
47 Kay, J. C., G. C. Claghorn, Z. Thompson, T. G. Hampton, and T. Garland, Jr. 2019.  
48  
49 Electrocardiograms of mice selectively bred for high levels of voluntary exercise:  
50  
51 Effects of short-term exercise training and the mini-muscle phenotype. *Physiol.*  
52  
53 *Behav.* 199:322–332.  
54  
55  
56  
57 Keeney, B. K., T. H. Meek, K. M. Middleton, L. F. Holness, and T. Garland, Jr. 2012. Sex  
58  
59 differences in cannabinoid receptor-1 (CB1) pharmacology in mice selectively bred  
60  
61  
62  
63  
64  
65

- for high voluntary wheel-running behavior. *Pharmacol. Biochem. Behav.* 101:528–537.
- Kelly, S. A., T. A. Bell, S. R. Selitsky, R. J. Buus, K. Hua, G. M. Weinstock, T. Garland, Jr., F. Pardo-Manuel de Villena, and D. Pomp. 2013. A novel intronic single nucleotide polymorphism in the *Myosin heavy polypeptide 4* gene is responsible for the Mini-Muscle phenotype characterized by major reduction in hind-limb muscle mass in mice. *Genetics* 195:1385–1395.
- Kelly, S. A., F. R. Gomes, E. M. Kolb, J. L. Malisch, and T. Garland, Jr. 2017. Effects of activity, genetic selection and their interaction on muscle metabolic capacities and organ masses in mice. *J. Exp. Biol.* 220:1038–1047.
- Kirk, S. F. L., T. L. Penney, and T.-L. F. McHugh. 2010. Characterizing the obesogenic environment: the state of the evidence with directions for future research. *Obes. Rev.* 11:109–117.
- Kirwan, J. P., J. Sacks, and S. Nieuwoudt. 2017. The essential role of exercise in the management of type 2 diabetes. *Cleve. Clin. J. Med.* 84:S15–S21.
- Klinker, F., K. Köhnemann, W. Paulus, and D. Liebetanz. 2017. Dopamine D3 receptor status modulates sexual dimorphism in voluntary wheel running behavior in mice. *Behav. Brain Res.* 333:235–241.
- Koch, L. G., and S. L. Britton. 2001. Artificial selection for intrinsic aerobic endurance running capacity in rats. *Physiol. Genomics* 5:45–52.
- Koeneman, M. A., M. W. Verheijden, M. J. M. Chinapaw, and M. Hopman-Rock. 2011. Determinants of physical activity and exercise in healthy older adults: a systematic review. *Int. J. Behav. Nutr. Phys. Act.* 8:142.

- Kolb, E. M., S. A. Kelly, K. M. Middleton, L. S. Sermsakdi, M. A. Chappell, and T. Garland, Jr. 2010. Erythropoietin elevates but not voluntary wheel running in mice. *J. Exp. Biol.* 213:510–519.
- Korte, S. M. 2001. Corticosteroids in relation to fear, anxiety and psychopathology. *Neurosci. Biobehav. Rev.* 25:117–142.
- Koteja, P., P. A. Carter, J. G. Swallow, and T. Garland, Jr. 2003. Food wasting by house mice: variation among individuals, families, and genetic lines. *Physiol. Behav.* 80:375–383.
- Langley-Evans, S. C. 2009. Nutritional programming of disease: unravelling the mechanism. *J. Anat.* 215:36–51.
- Lee, B. Y., S. M. Bartsch, Y. Mui, L. A. Haidari, M. L. Spiker, and J. Gittelsohn. 2017. A systems approach to obesity. *Nutr. Rev.* 75:94–106.
- Leszczynski, E. C., J. R. Visker, and D. P. Ferguson. 2019. The effect of growth restriction on voluntary physical activity engagement in mice. *Med. Sci. Sports Exerc.* 51:2201–2209.
- Levine, J. A. 2007. Nonexercise activity thermogenesis--liberating the life-force. *J. Intern. Med.* 262:273–287.
- Levine, J. A., S. K. McCrady, L. M. Lanningham-Foster, P. H. Kane, R. C. Foster, and C. U. Manohar. 2008. The role of free-living daily walking in human weight gain and obesity. *Diabetes* 57:548–554.
- Li, G., J. J. Kohorst, W. Zhang, E. Laritsky, G. Kunde-Ramamoorthy, M. S. Baker, M. L. Fiorotto, and R. A. Waterland. 2013. Early postnatal nutrition determines adult physical activity and energy expenditure in female mice. *Diabetes* 62:2773–2783.



- 1  
2  
3  
4 Lightfoot, J. T. 2008. Sex hormones' regulation of rodent physical activity: a review. *Int. J.*  
5  
6 *Biol. Sci.* 4:126–132.  
7  
8  
9 Lightfoot, J. T., E. J. C. DE Geus, F. W. Booth, M. S. Bray, M. DEN Hoed, J. Kaprio, S. A.  
10  
11 Kelly, D. Pomp, M. C. Saul, M. A. Thomis, T. Garland, Jr., and C. Bouchard. 2018.  
12  
13 Biological/Genetic Regulation of Physical Activity Level: Consensus from  
14  
15 GenBioPAC. *Med. Sci. Sports Exerc.* 50:863–873.  
16  
17  
18 Lloyd, L. J., S. C. Langley-Evans, and S. McMullen. 2012. Childhood obesity and risk of  
19  
20 the adult metabolic syndrome: a systematic review. *Int. J. Obes.* 36:1–11.  
21  
22  
23 Lussana, F., R. C. Painter, M. C. Ocke, H. R. Buller, P. M. Bossuyt, and T. J. Roseboom.  
24  
25 2008. Prenatal exposure to the Dutch famine is associated with a preference for  
26  
27 fatty foods and a more atherogenic lipid profile. *Am. J. Clin. Nutr.* 88:1648–1652.  
28  
29  
30 Malina, R. M. 2001. Tracking of physical activity across the lifespan.  
31  
32  
33 Malisch, J. L., C. W. Breuner, F. R. Gomes, M. A. Chappell, and T. Garland, Jr. 2008.  
34  
35 Circadian pattern of total and free corticosterone concentrations, corticosteroid-  
36  
37 binding globulin, and physical activity in mice selectively bred for high voluntary  
38  
39 wheel-running behavior. *Gen. Comp. Endocrinol.* 156:210–217.  
40  
41  
42 Malisch, J. L., K. deWolski, T. H. Meek, W. Acosta, K. M. Middleton, O. L. Crino, and T.  
43  
44 Garland, Jr. 2016. Acute restraint stress alters wheel-running behavior  
45  
46 immediately following stress and up to 20 hours later in house mice. *Physiol.*  
47  
48 *Biochem. Zool.* 89:546–552.  
49  
50  
51 Malisch, J. L., W. Saltzman, F. R. Gomes, E. L. Rezende, D. R. Jeske, and T. Garland,  
52  
53 Jr. 2007. Baseline and stress-induced plasma corticosterone concentrations of  
54  
55 mice selectively bred for high voluntary wheel running. *Physiol. Biochem. Zool.*  
56  
57  
58 80:146–156.  
59  
60  
61  
62  
63  
64  
65

- 1  
2  
3  
4 Mattioli, A. V., S. Sciomer, C. Cocchi, S. Maffei, and S. Gallina. 2020. Quarantine during  
5  
6 COVID-19 outbreak: Changes in diet and physical activity increase the risk of  
7  
8 cardiovascular disease. *Nutr. Metab. Cardiovasc. Dis.* 30:1409–1417.  
9
- 10  
11 Maurice, J., and P. Manousou. 2018. Non-alcoholic fatty liver disease. *Clin. Med. Lond.*  
12  
13 *Engl.* 18:245–250.  
14
- 15  
16 McAllister, E. J., N. V. Dhurandhar, S. W. Keith, L. J. Aronne, J. Barger, M. Baskin, R. M.  
17  
18 Benca, J. Biggio, M. M. Boggiano, J. C. Eisenmann, M. Elobeid, K. R. Fontaine, P.  
19  
20 Gluckman, E. C. Hanlon, P. Katzmarzyk, A. Pietrobelli, D. T. Redden, D. M.  
21  
22 Ruden, C. Wang, R. A. Waterland, S. M. Wright, and D. B. Allison. 2009. Ten  
23  
24 Putative Contributors to the Obesity Epidemic. *Crit. Rev. Food Sci. Nutr.* 49:868–  
25  
26 913.  
27  
28  
29
- 30  
31 McEwen, B. S., J. D. Gray, and C. Nasca. 2015. 60 YEARS OF  
32  
33 NEUROENDOCRINOLOGY: Redefining neuroendocrinology: stress, sex and  
34  
35 cognitive and emotional regulation. *J. Endocrinol.* 226:T67–T83.  
36  
37
- 38  
39 McGuire, K. A., and R. Ross. 2011. Incidental physical activity is positively associated  
40  
41 with cardiorespiratory fitness. *Med. Sci. Sports Exerc.* 43:2189–2194.  
42
- 43  
44 McLeod, M., L. Breen, D. L. Hamilton, and A. Philp. 2016. Live strong and prosper: the  
45  
46 importance of skeletal muscle strength for healthy ageing. *Biogerontology* 17:497–  
47  
48 510.  
49
- 50  
51 McNamara, M. P., J. M. Singleton, M. D. Cadney, P. M. Ruegger, J. Borneman, and T.  
52  
53 Garland, Jr. 2021. Early-life effects of juvenile Western diet and exercise on adult  
54  
55 gut microbiome composition in mice. *J. Exp. Biol.* (In press):jeb.239699.  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2  
3 Meek, T. H., E. M. Dlugosz, K. T. Vu, and T. Garland, Jr. 2012. Effects of leptin treatment  
4 and Western diet on wheel running in selectively bred high runner mice. *Physiol.*  
5  
6 *Behav.* 106:252–258.  
7  
8  
9
- 10 Meek, T. H., J. C. Eisenmann, and T. Garland, Jr. 2010. Western diet increases wheel  
11 running in mice selectively bred for high voluntary wheel running. *Int. J. Obes.*  
12  
13 34:960–969.  
14  
15  
16
- 17 Meek, T. H., J. C. Eisenmann, B. K. Keeney, R. M. Hannon, E. M. Dlugosz, and T.  
18 Garland, Jr. 2014. Effects of early-life exposure to Western diet and wheel access  
19 on metabolic syndrome profiles in mice bred for high voluntary exercise. *Genes*  
20 *Brain Behav.* 13:322–332.  
21  
22  
23
- 24 Meek, T. H., and G. J. Morton. 2016. The role of leptin in diabetes: metabolic effects.  
25  
26 *Diabetologia* 59:928–932.  
27  
28  
29
- 30 Metcalf, B. S., J. Hosking, A. N. Jeffery, L. D. Voss, W. Henley, and T. J. Wilkin. 2011.  
31  
32 Fatness leads to inactivity, but inactivity does not lead to fatness: a longitudinal  
33 study in children (EarlyBird 45). *Arch. Dis. Child.* 96:942–947.  
34  
35  
36
- 37 Metges, C. C. 2009. Early nutrition and later obesity: animal models provide insights into  
38 mechanisms. *Adv. Exp. Med. Biol.* 646:105–112.  
39  
40  
41
- 42 Mika, A., W. V. Treuren, A. González, J. J. Herrera, R. Knight, and M. Fleshner. 2015.  
43  
44 Exercise is more effective at altering gut microbial composition and producing  
45 stable changes in lean mass in juvenile versus adult male F344 rats. *PLOS ONE*  
46  
47 10:e0125889.  
48  
49  
50
- 51 Miles, J. L., K. Huber, N. M. Thompson, M. Davison, and B. H. Breier. 2009. Moderate  
52 daily exercise activates metabolic flexibility to prevent prenatally induced obesity.  
53  
54 *Endocrinology* 150:179–186.  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2  
3 Mokhtari-Zaer, A., M. Hosseini, and M. H. Boskabady. 2018. The effects of exercise on  
4  
5 depressive- and anxiety-like behaviors as well as lung and hippocampus oxidative  
6  
7 stress in ovalbumin-sensitized juvenile rats. *Respir. Physiol. Neurobiol.* 248:55–62.  
8  
9
- 10 Moraal, M., P. P. a. M. Leenaars, H. Arnts, K. Smeets, B. S. Savenije, J. H. a. J. Curfs,  
11  
12 and M. Ritskes-Hoitinga. 2012. The influence of food restriction versus ad libitum  
13  
14 feeding of chow and purified diets on variation in body weight, growth and  
15  
16 physiology of female Wistar rats. *Lab. Anim.* 46:101–107.  
17  
18
- 19 Morrison, J. A., L. A. Friedman, and C. Gray-McGuire. 2007. Metabolic syndrome in  
20  
21 childhood predicts adult cardiovascular disease 25 years later: the Princeton Lipid  
22  
23 Research Clinics Follow-up Study. *Pediatrics* 120:340–345.  
24  
25
- 26 Nathanielsz, P. W. 2006. Animal models that elucidate basic principles of the  
27  
28 developmental origins of adult diseases. *ILAR J.* 47:73–82.  
29  
30
- 31 Negaresh, R., R. Motl, J. Baker, P. Zimmer, N. Majdinasab, D. Patel, M. Khodadoost, R.  
32  
33 Ranjbar, and M. Mokhtarzade. 2018. Exercise-induced changes in neurotrophic  
34  
35 factors and markers of blood-brain barrier permeability are moderated by weight  
36  
37 status in multiple sclerosis. *Neuropeptides* 70:93–100.  
38  
39
- 40 Noland, R. C., J. P. Thyfault, S. T. Henes, B. R. Whitfield, T. L. Woodlief, J. R. Evans, J.  
41  
42 A. Lust, S. L. Britton, L. G. Koch, R. W. Dudek, G. L. Dohm, R. N. Cortright, and R.  
43  
44 M. Lust. 2007. Artificial selection for high-capacity endurance running is protective  
45  
46 against high-fat diet-induced insulin resistance. *Am. J. Physiol. Endocrinol. Metab.*  
47  
48 293:E31-41.  
49  
50
- 51 North, K. E., M. Graff, L. S. Adair, E. M. Lange, L. A. Lange, G. Guo, and P. Gordon-  
52  
53 Larsen. 2010. Genetic epidemiology of body mass index and body mass change  
54  
55 from adolescence to young adulthood. *Obes. Silver Spring Md* 18:1474–1476.  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- Novak, C. M., C. Escande, S. M. Gerber, E. N. Chini, M. Zhang, S. L. Britton, L. G. Koch, and J. A. Levine. 2009. Endurance capacity, not body size, determines physical activity levels: role of skeletal muscle PEPCK. *PloS One* 4:e5869.
- Obert, J., M. Pearlman, L. Obert, and S. Chapin. 2017. Popular weight loss strategies: a review of four weight loss techniques. *Curr. Gastroenterol. Rep.* 19:61.
- Oelkrug, R., C. Krause, B. Herrmann, J. Resch, S. Gachkar, A. T. El Gammal, S. Wolter, O. Mann, H. Oster, H. Kirchner, and J. Mittag. 2020. Maternal brown fat thermogenesis programs glucose tolerance in the male offspring. *Cell Rep.* 33:108351.
- O'Neill, B., and P. Raggi. 2020. The ketogenic diet: Pros and cons. *Atherosclerosis* 292:119–126.
- Orsso, C. E., J. R. B. Tibaes, C. L. P. Oliveira, D. A. Rubin, C. J. Field, S. B. Heymsfield, C. M. Prado, and A. M. Haqq. 2019. Low muscle mass and strength in pediatrics patients: Why should we care? *Clin. Nutr.* 38:2002–2015.
- Ozanne, S. E., and K. Siddle. 2011. Developmental origins of insulin resistance and type 2 diabetes. Pp. 60–80 *in* The metabolic syndrome. John Wiley & Sons, Ltd.
- Paglia, L. 2019. The sweet danger of added sugars. *Eur. J. Paediatr. Dent.* 20:89.
- Palmer, B. F., and D. J. Clegg. 2015. The sexual dimorphism of obesity. *Mol. Cell. Endocrinol.* 402:113–119.
- Pan, Y., and C. A. Pratt. 2008. Metabolic syndrome and its association with diet and physical activity in US adolescents. *J. Am. Diet. Assoc.* 108:276–286; discussion 286.
- Papas, M. A., A. J. Alberg, R. Ewing, K. J. Helzlsouer, T. L. Gary, and A. C. Klassen. 2007. The built environment and obesity. *Epidemiol. Rev.* 29:129–143.

- 1  
2  
3  
4 Park, J.-S., and A. Höke. 2014. Treadmill exercise induced functional recovery after  
5  
6 peripheral nerve repair is associated with increased levels of neurotrophic factors.  
7  
8 PLOS ONE 9:e90245.  
9
- 10  
11 Patterson, C. M., S. G. Bouret, A. A. Dunn-Meynell, and B. E. Levin. 2009. Three weeks  
12  
13 of postweaning exercise in DIO rats produces prolonged increases in central leptin  
14  
15 sensitivity and signaling. *Am. J. Physiol. - Regul. Integr. Comp. Physiol.*  
16  
17 296:R537–R548.  
18  
19
- 20  
21 Patterson, C. M., A. A. Dunn-Meynell, and B. E. Levin. 2008. Three weeks of early-onset  
22  
23 exercise prolongs obesity resistance in DIO rats after exercise cessation. *Am. J.*  
24  
25 *Physiol. Regul. Integr. Comp. Physiol.* 294:R290-301.  
26  
27
- 28  
29 Patterson, Z. R., and A. B. Abizaid. 2013. Stress induced obesity: lessons from rodent  
30  
31 models of stress. *Front. Neurosci.* 7:130.  
32
- 33  
34 Peters, L. L., R. F. Robledo, C. J. Bult, G. A. Churchill, B. J. Paigen, and K. L. Svenson.  
35  
36 2007. The mouse as a model for human biology: a resource guide for complex trait  
37  
38 analysis. *Nat. Rev. Genet.* 8:58–69. Nature Publishing Group.  
39
- 40  
41 Rankinen, T., and C. Bouchard. 2008. Gene–physical activity interactions: overview of  
42  
43 human studies. *Obes. Silver Spring Md* 16:S47–S50.  
44
- 45  
46 Reilly, J. J. 2011. Can we modulate physical activity in children? *Int. J. Obes.* 35:1266–  
47  
48 1269. Nature Publishing Group.  
49
- 50  
51 Reilly, J. J. 2016. When does it all go wrong? Longitudinal studies of changes in  
52  
53 moderate-to-vigorous-intensity physical activity across childhood  
54  
55 and adolescence. *J. Exerc. Sci. Fit.* 14:1–6.  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2  
3  
4 Rezende, E. L. 2006. Maximum aerobic performance in lines of Mus selected for high  
5  
6 wheel-running activity: effects of selection, oxygen availability and the mini-muscle  
7  
8 phenotype. J. Exp. Biol. 209:115–127.  
9
- 10  
11 Rhodes, J. S., S. C. Gammie, and T. Garland, Jr. 2005. Neurobiology of mice selected  
12  
13 for high voluntary wheel-running activity. Integr. Comp. Biol. 45:438–455.  
14
- 15  
16 Rhodes, J. S., and T. J. Kawecki. 2009. Behavior and neurobiology. Pp. 263–300 in T.  
17  
18 Garland, Jr. and M. R. Rose, eds. Experimental evolution: concepts, methods, and  
19  
20 applications of selection experiments. University of California Press, Berkeley.  
21  
22
- 23 Roberto, C. A., B. Swinburn, C. Hawkes, T. T.-K. Huang, S. A. Costa, M. Ashe, L.  
24  
25 Zwicker, J. H. Cawley, and K. D. Brownell. 2015. Patchy progress on obesity  
26  
27 prevention: emerging examples, entrenched barriers, and new thinking. The  
28  
29 Lancet 385:2400–2409.  
30  
31
- 32  
33 Roberts, M. D., L. Gilpin, K. E. Parker, T. E. Childs, M. J. Will, and F. W. Booth. 2012.  
34  
35 Dopamine D1 receptor modulation in nucleus accumbens lowers voluntary wheel  
36  
37 running in rats bred to run high distances. Physiol. Behav. 105:661–668.  
38  
39
- 40 Roghair, R. D., J. L. Segar, K. A. Volk, M. W. Chapleau, L. M. Dallas, A. R. Sorenson, T.  
41  
42 D. Scholz, and F. S. Lamb. 2009. Vascular nitric oxide and superoxide anion  
43  
44 contribute to sex-specific programmed cardiovascular physiology in mice. Am. J.  
45  
46 Physiol. - Regul. Integr. Comp. Physiol. 296:R651–R662.  
47  
48
- 49  
50 Rohde, K., M. Keller, L. la C. Poulsen, M. Blüher, P. Kovacs, and Y. Böttcher. 2019.  
51  
52 Genetics and epigenetics in obesity. Metab. - Clin. Exp. 92:37–50.  
53  
54
- 55 Rosa, B. V., H. T. Blair, M. H. Vickers, K. E. Dittmer, P. C. H. Morel, C. G. Knight, and E.  
56  
57 C. Firth. 2013. Moderate exercise during pregnancy in wistar rats alters bone and  
58  
59  
60  
61  
62  
63  
64  
65

- body composition of the adult offspring in a sex-dependent manner. PLoS ONE 8:e82378.
- Rosenfeld, C. S. 2017. Sex-dependent differences in voluntary physical activity. J. Neurosci. Res. 95:279–290.
- Ross, M. G., and M. Desai. 2005. Gestational programming: population survival effects of drought and famine during pregnancy. Am. J. Physiol. Regul. Integr. Comp. Physiol. 288:R25-33.
- Rowland, T. W. 1998. The biological basis of physical activity. Med. Sci. Sports Exerc. 30:392–399.
- Ruiz-Roso, M. B., P. de Carvalho Padilha, D. C. Mantilla-Escalante, N. Ulloa, P. Brun, D. Acevedo-Correa, W. Arantes Ferreira Peres, M. Martorell, M. T. Aires, L. de Oliveira Cardoso, F. Carrasco-Marín, K. Paternina-Sierra, J. E. Rodriguez-Meza, P. M. Montero, G. Bernabè, A. Pauletto, X. Taci, F. Visioli, and A. Dávalos. 2020. COVID-19 confinement and changes of adolescent's dietary trends in Italy, Spain, Chile, Colombia and Brazil. Nutrients 12.
- Rundle, A. G., Y. Park, J. B. Herbstman, E. W. Kinsey, and Y. C. Wang. 2020. COVID-19–related school closings and risk of weight gain among children. Obesity 28:1008–1009.
- Ryan, D. H., and Diabetes Prevention Program Research Group. 2003. Diet and exercise in the prevention of diabetes. Int. J. Clin. Pract. Suppl. (134):28–35.
- Sallis, J. F., and M. F. Hovell. 1990. Determinants of exercise behavior. Exerc. Sport Sci. Rev. 18:307–330.
- Samuelsson, A.-M., P. A. Matthews, M. Argenton, M. R. Christie, J. M. McConnell, E. H. J. M. Jansen, A. H. Piersma, S. E. Ozanne, D. F. Twinn, C. Remacle, A.



- Rowlerson, L. Poston, and P. D. Taylor. 2008. Diet-induced obesity in female mice leads to offspring hyperphagia, adiposity, hypertension, and insulin resistance: a novel murine model of developmental programming. *Hypertension* 51:383–392.
- Samuelsson, A.-M., A. Morris, N. Igosheva, S. L. Kirk, J. M. C. Pombo, C. W. Coen, L. Poston, and P. D. Taylor. 2010. Evidence for sympathetic origins of hypertension in juvenile offspring of obese rats. *Hypertension* 55:76–82.
- Schroeder, M., L. Shbiro, V. Gelber, and A. Weller. 2010. Post-weaning voluntary exercise exerts long-term moderation of adiposity in males but not in females in an animal model of early-onset obesity. *Horm. Behav.* 57:496–505.
- Sidor, A., and P. Rzymiski. 2020. Dietary choices and habits during COVID-19 lockdown: experience from poland. *Nutrients* 12:13.
- Simi, B., B. Sempore, M. H. Mayet, and R. J. Favier. 1991. Additive effects of training and high-fat diet on energy metabolism during exercise. *J. Appl. Physiol.* 71:197–203.
- Simmons, R. A. 2009. Developmental origins of adult disease. *Pediatr. Clin. North Am.* 56:449–466.
- Singh, A. S., C. Mulder, J. W. R. Twisk, W. van Mechelen, and M. J. M. Chinapaw. 2008. Tracking of childhood overweight into adulthood: a systematic review of the literature. *Obes. Rev. Off. J. Int. Assoc. Study Obes.* 9:474–488.
- Sisson, S. B., S. M. Camhi, T. S. Church, C. K. Martin, C. Tudor-Locke, C. Bouchard, C. P. Earnest, S. R. Smith, R. L. Newton, T. Rankinen, and P. T. Katzmarzyk. 2009. Leisure time sedentary behavior, occupational/domestic physical activity, and metabolic syndrome in U.S. men and women. *Metab. Syndr. Relat. Disord.* 7:529–536.

Stanford, K. I., M.-Y. Lee, K. M. Getchell, K. So, M. F. Hirshman, and L. J. Goodyear.

2015. Exercise before and during pregnancy prevents the deleterious effects of maternal high-fat feeding on metabolic health of male offspring. *Diabetes* 64:427–433.

Stanton, R., Q. G. To, S. Khalesi, S. L. Williams, S. J. Alley, T. L. Thwaite, A. S. Fenning, and C. Vandelanotte. 2020. Depression, anxiety and stress during COVID-19: associations with changes in physical activity, sleep, tobacco and alcohol use in Australian adults. *Int. J. Environ. Res. Public. Health* 17:13.

Stein, A. D., A. Rundle, N. Wada, R. A. Goldbohm, and L. H. Lumey. 2009. Associations of gestational exposure to famine with energy balance and macronutrient density of the diet at age 58 years differ according to the reference population used. *J. Nutr.* 139:1555–1561.

Steinberger, J., S. R. Daniels, R. H. Eckel, L. Hayman, R. H. Lustig, B. McCrindle, M. L. Mietus-Snyder, and American Heart Association Atherosclerosis, Hypertension, and Obesity in the Young Committee of the Council on Cardiovascular Disease in the Young; Council on Cardiovascular Nursing; and Council on Nutrition, Physical Activity, and Metabolism. 2009. Progress and challenges in metabolic syndrome in children and adolescents: a scientific statement from the American Heart Association Atherosclerosis, Hypertension, and Obesity in the Young Committee of the Council on Cardiovascular Disease in the Young; Council on Cardiovascular Nursing; and Council on Nutrition, Physical Activity, and Metabolism. *Circulation* 119:628–647.

Strohacker, K., K. C. Carpenter, and B. K. McFarlin. 2009. Consequences of weight cycling: an increase in disease risk? *Int. J. Exerc. Sci.* 2:191–201.

- 1  
2  
3  
4 Strohacker, K., O. Galarraga, and D. M. Williams. 2014. The Impact of Incentives on  
5  
6 Exercise Behavior: A Systematic Review of Randomized Controlled Trials. *Ann.*  
7  
8 *Behav. Med. Publ. Soc. Behav. Med.* 48:92–99.  
9
- 10  
11 Sumiyoshi, A., Y. Taki, H. Nonaka, H. Takeuchi, and R. Kawashima. 2014. Regional gray  
12  
13 matter volume increases following 7 days of voluntary wheel running exercise: A  
14  
15 longitudinal VBM study in rats. *NeuroImage* 98:82–90.  
16  
17
- 18  
19 Sun, B., N.-C. Liang, E. R. Ewald, R. H. Purcell, G. J. Boersma, J. Yan, T. H. Moran, and  
20  
21 K. L. K. Tamashiro. 2013. Early postweaning exercise improves central leptin  
22  
23 sensitivity in offspring of rat dams fed high-fat diet during pregnancy and lactation.  
24  
25 *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 305:R1076-1084.  
26  
27
- 28  
29 Sutton, G. M., A. V. Centanni, and A. A. Butler. 2010. Protein malnutrition during  
30  
31 pregnancy in C57BL/6J mice results in offspring with altered circadian physiology  
32  
33 before obesity. *Endocrinology* 151:1570–1580.  
34
- 35  
36 Svenson, K. L., R. Von Smith, P. A. Magnani, H. R. Suetin, B. Paigen, J. K. Naggert, R.  
37  
38 Li, G. A. Churchill, and L. L. Peters. 2007. Multiple trait measurements in 43 inbred  
39  
40 mouse strains capture the phenotypic diversity characteristic of human  
41  
42 populations. *J. Appl. Physiol.* 102:2369–2378.  
43  
44
- 45  
46 Swallow, J. G., P. A. Carter, and T. Garland. 1998. Artificial selection for increased  
47  
48 wheel-running behavior in house mice. *Behav. Genet.* 28:227–237.  
49
- 50  
51 Swallow, J. G., and T. Garland, Jr. 2005. Selection experiments as a tool in evolutionary  
52  
53 and comparative physiology: insights into complex traits--an introduction to the  
54  
55 symposium. *Integr. Comp. Biol.* 45:387–390.  
56
- 57  
58 Swallow, J. G., J. P. Hayes, P. Koteja, and T. Garland, Jr. 2009. Selection experiments  
59  
60 and experimental evolution of performance and physiology. Pp. 301–351 *in*  
61  
62  
63  
64  
65

- Theodore Garland, Jr. and Michael R. Rose, eds. Experimental evolution: concepts, methods, and applications of selection experiments. University of California Press, Berkeley.
- Swallow, J. G., P. Koteja, P. A. Carter, and T. Garland, Jr. 1999. Artificial selection for increased wheel-running activity in house mice results in decreased body mass at maturity. *J. Exp. Biol.* 202:2513–2520.
- Swallow, J., P. Koteja, P. Carter, and T. Garland, Jr. 2001. Food consumption and body composition in mice selected for high wheel-running activity. *J. Comp. Physiol. B* 171:651–659.
- Swift, D. L., N. M. Johannsen, C. J. Lavie, C. P. Earnest, and T. S. Church. 2014. The role of exercise and physical activity in weight loss and maintenance. *Prog. Cardiovasc. Dis.* 56:441–447.
- Syed, S. A., E. L. P. Smith, O. Batuman, L. A. Rosenblum, M. J. Owens, C. B. Nemeroff, and J. D. Coplan. 2017. Adult glucocorticoid receptor mRNA expression volatility in response to an acute stressor and juvenile CSF corticotropin-releasing factor: A pilot neurodevelopmental study. *Neurosci. Lett.* 647:20–25.
- Syme, D. A., K. Evashuk, B. Grintuch, E. L. Rezende, and T. Garland, Jr. 2005. Contractile abilities of normal and “mini” triceps surae muscles from mice (*Mus domesticus*) selectively bred for high voluntary wheel running. *J. Appl. Physiol.* 99:1308–1316.
- Talmadge, R. J., W. Acosta, and T. Garland, Jr. 2014. Myosin heavy chain isoform expression in adult and juvenile mini-muscle mice bred for high-voluntary wheel running. *Mech. Dev.* 134:16–30.

- 1  
2  
3  
4 Teegarden, S. L., A. N. Scott, and T. L. Bale. 2009. Early life exposure to a high fat diet  
5  
6 promotes long-term changes in dietary preferences and central reward signaling.  
7  
8 Neuroscience 162:924–932.  
9
- 10  
11 Telama, R., X. Yang, J. Viikari, I. Välimäki, O. Wanne, and O. Raitakari. 2005. Physical  
12  
13 activity from childhood to adulthood. Am. J. Prev. Med. 28:267–273.  
14
- 15  
16 Teske, J. A., C. J. Billington, and C. M. Kotz. 2014. Mechanisms underlying obesity  
17  
18 resistance associated with high spontaneous physical activity. Neuroscience  
19  
20 256:91–100.  
21  
22
- 23  
24 The, N. S., C. Suchindran, K. E. North, B. M. Popkin, and P. Gordon-Larsen. 2010.  
25  
26 Association of adolescent obesity with risk of severe obesity in adulthood. JAMA  
27  
28 304:2042–2047.  
29
- 30  
31 Thompson, Z., D. Argueta, T. Garland, Jr., and N. DiPatrizio. 2017. Circulating levels of  
32  
33 endocannabinoids respond acutely to voluntary exercise, are altered in mice  
34  
35 selectively bred for high voluntary wheel running, and differ between the sexes.  
36  
37 Physiol. Behav. 170:141–150.  
38  
39
- 40  
41 Triantafyllou, G. A., S. A. Paschou, and C. S. Mantzoros. 2016. Leptin and hormones:  
42  
43 energy homeostasis. Endocrinol. Metab. Clin. North Am. 45:633–645.  
44
- 45  
46 Trost, S., N. Owen, A. Bauman, J. Sallis, and W. Brown. 2002. Correlates of adults'  
47  
48 participation in physical activity: review and update. Med. Sci. Sports Exerc.  
49  
50 34:1996–2001.  
51
- 52  
53 Twisk, J. W. 2001. Physical activity guidelines for children and adolescents: a critical  
54  
55 review. Sports Med. 31:617–627.  
56
- 57  
58 Vaanholt, L. M., I. Jonas, M. Doornbos, K. A. Schubert, C. Nyakas, T. Garland, Jr., G. H.  
59  
60 Visser, and G. van Dijk. 2008. Metabolic and behavioral responses to high-fat  
61  
62  
63  
64  
65


- feeding in mice selectively bred for high wheel-running activity. *Int. J. Obes.* 32:1566–1575.
- Vaanholt, L., P. Meerlo, T. Garland, Jr., G. Visser, and G. van Dijk. 2007. Plasma adiponectin is increased in mice selectively bred for high wheel-running activity, but not by wheel running per sé. *Horm. Metab. Res.* 39:377–383.
- van Keulen, B. J., C. V. Dolan, B. van der Voorn, R. Andrew, B. R. Walker, H. Hulshoff Pol, D. I. Boomsma, J. Rotteveel, and M. J. J. Finken. 2020. Sexual dimorphism in cortisol metabolism throughout pubertal development: a longitudinal study. *Endocr. Connect.* 9:542–551.
- Vickers, M. H., B. H. Breier, D. McCarthy, and P. D. Gluckman. 2003. Sedentary behavior during postnatal life is determined by the prenatal environment and exacerbated by postnatal hypercaloric nutrition. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 285:R271-273.
- Vucetic, Z., J. L. Carlin, K. Totoki, and T. M. Reyes. 2012. Epigenetic dysregulation of the dopamine system in diet-induced obesity. *J. Neurochem.* 120:891–898.
- Vucetic, Z., J. Kimmel, K. Totoki, E. Hollenbeck, and T. M. Reyes. 2010. Maternal high-fat diet alters methylation and gene expression of dopamine and opioid-related genes. *Endocrinology* 151:4756–4764.
- Wallace, I. J., and T. Garland, Jr. 2016. Mobility as an emergent property of biological organization: insights from experimental evolution. *Evol. Anthropol.* 25:98–104.
- Wang, G., Y. Zhang, J. Zhao, J. Zhang, and F. Jiang. 2020. Mitigate the effects of home confinement on children during the COVID-19 outbreak. *The Lancet* 395:945–947.

- 1  
2  
3  
4 Warden, S. J., R. K. Fuchs, A. B. Castillo, I. R. Nelson, and C. H. Turner. 2007. Exercise  
5  
6 when young provides lifelong benefits to bone structure and strength. *J. Bone*  
7  
8 *Miner. Res.* 22:251–259.  
9
- 10  
11 Wells, J. C. K. 2007. Sexual dimorphism of body composition. *Best Pract. Res. Clin.*  
12  
13 *Endocrinol. Metab.* 21:415–430.  
14
- 15  
16 Westcott, W. L. 2012. Resistance training is medicine: effects of strength training on  
17  
18 health. *Curr. Sports Med. Rep.* 11:209–216.  
19
- 20  
21 Westerterp, K. R., and J. R. Speakman. 2008. Physical activity energy expenditure has  
22  
23 not declined since the 1980s and matches energy expenditures of wild mammals.  
24  
25 *Int. J. Obes.* 2005 32:1256–1263.  
26
- 27  
28 Whitaker, K. W., K. Totoki, and T. M. Reyes. 2012. Metabolic adaptations to early life  
29  
30 protein restriction differ by offspring sex and postweaning diet in the mouse. *Nutr.*  
31  
32 *Metab. Cardiovasc. Dis.* 22:1067–1074.  
33  
34
- 35  
36 Wilkin, T. J. 2011. Can we modulate physical activity in children? No. *Int. J. Obes.* 2005  
37  
38 35:1270–1276.  
39
- 40  
41 Woods, J. A., N. T. Hutchinson, S. K. Powers, W. O. Roberts, M. C. Gomez-Cabrera, Z.  
42  
43 Radak, I. Berkes, A. Boros, I. Boldogh, C. Leeuwenburgh, H. J. Coelho-Júnior, E.  
44  
45 Marzetti, Y. Cheng, J. Liu, J. L. Durstine, J. Sun, and L. L. Ji. 2020. The COVID-19  
46  
47 pandemic and physical activity. *Sports Med. Health Sci.* 2:55–64.  
48  
49
- 50  
51 Yang, Z., and S. L. Huffman. 2013. Nutrition in pregnancy and early childhood and  
52  
53 associations with obesity in developing countries. *Matern. Child. Nutr.* 9 Suppl  
54  
55 1:105–119.  
56
- 57  
58 Zhang, F., Y. Chen, M. Heiman, and R. Dimarchi. 2005. Leptin: structure, function and  
59  
60 biology. *Vitam. Horm.* 71:345–372.  
61  
62  
63  
64  
65


- 1  
2  
3  
4 Zhang, R., M. Asai, C. E. Mahoney, M. Joachim, Y. Shen, G. Gunner, and J. A. Majzoub.  
5  
6 2017. Loss of hypothalamic corticotropin-releasing hormone markedly reduces  
7  
8 anxiety behaviors in mice. *Mol. Psychiatry* 22:733–744.  
9
- 10  
11 Zhang, Y., and S. Chua. 2017. Leptin function and regulation. *Compr. Physiol.* 8:351–  
12  
13 369.  
14
- 15  
16 Zhou, W.-C., Q.-B. Zhang, and L. Qiao. 2014. Pathogenesis of liver cirrhosis. *World J.*  
17  
18 *Gastroenterol.* 20:7312–7324.  
19
- 20  
21 Zombeck, J. A., E. K. Deyoung, W. J. Brzezinska, and J. S. Rhodes. 2011. Selective  
22  
23 breeding for increased home cage physical activity in collaborative cross and  
24  
25 Hsd:ICR mice. *Behav. Genet.* 41:571–582.  
26
- 27  
28 Zore, T., M. Palafox, and K. Reue. 2018. Sex differences in obesity, lipid metabolism, and  
29  
30 inflammation-A role for the sex chromosomes? *Mol. Metab.* 15:35–44.  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65




**Figure 1. Experimental timeline** of events starting with the first births of generation 76 mice. Of these mice, 192 males were used for the present experiment. Three-week-long early-life diet and exercise manipulation began immediately after weaning and was followed by an 8-week washout period, during which mice were individually housed without wheels and given standard diet. At the end of washout, all mice were tested. Note that corticosterone was measured in plasma taken before wheel running, whereas leptin and adiponectin were measured in terminal blood samples collected at dissections. WD = Western diet. Home-cage activity is an indicator of spontaneous physical activity (SPA).




**Figure 2. Juvenile wheel running** of male mice during the 3-week treatment period, shown as least squares means and standard errors with accompanying p-values from SAS Procedure Mixed. Note the data shown are from the experimental group given access to wheels (see Fig. 1). A) Mean wheel revolutions per day (circumference 1.12 meters), B) duration of daily running, C) mean revolutions per minute, D) maximum revolutions per minute. Above each week (set of four bars) of measurements are tables of main and interactive effects ( $p < 0.05$  in bold font). See Supplemental Table 1 for additional statistical details.




**Figure 3. Adult wheel running** during days 1-7 of a 2-week testing period. Values are least squares means and standard errors and accompanying p-values from SAS Procedure Mixed. A) Mean wheel revolutions per day (circumference 1.12 meters), B) duration of daily running, C) mean revolutions per minute, D) maximum revolutions per minute. Above each measurement are tables of main and interactive effects ( $p < 0.05$  in bold font). SD = standard diet; WD = Western diet; E = exercise; D = diet; LT = linetype. Values for the second week of adult testing can be seen in Supplemental Table 1.



**Figure 4. Total body mass, lean mass, and fat mass.** Values are least squares means and standard errors from SAS Procedure Mixed. Note that body composition (lean and fat mass) was not measured during weaning (week 3); however, HR mice at weaning weighed 12.61% less than C mice ( $p = 0.0278$ ). Fat mass was analyzed using lean mass as a covariate. See Supplemental Table 1 for additional statistical details. SD = standard diet, WD = Western diet, Sedentary = housed without wheels as juveniles, Exercise = housed with wheels as juveniles.



**Figure 5. Weekly mass-adjusted juvenile caloric intake** in response to juvenile diet and/or exercise treatment. Values are least squares means and standard errors and accompanying p-values from SAS Procedure Mixed. Above each week are tables of main and interactive effects (p < 0.05 in bold font). See Supplemental Table 1 for additional statistical details.



**Figure 6. Adult caloric intake** with body mass as a covariate. Values are least squares means and standard errors and accompanying p-values from SAS Procedure Mixed. Above each week are tables of main and interactive effects ( $p < 0.05$  in bold font). See Supplemental Table 1 for additional statistical details.




Figure 7. Maximal oxygen consumption ( $VO_2\text{max}$ ) measured during forced exercise (see Methods 2.3.4). Body mass was a significant predictor of  $VO_2\text{max}$  ( $p < 0.0001$ ), mice from High Runner lines had higher  $VO_2\text{max}$  than those from non-selected Control lines ( $p = 0.0019$ ), and individuals with the mini-muscle phenotype (not shown with a separate symbol) had higher values than normal-muscle mice ( $p = 0.0271$ ), with no effect of either early-life treatment and no interactions (see Supplemental Table 1 for additional statistical details).





Figure 8. Body-mass adjusted heart ventricles, triceps surae, brain, liver, and subdermal fat pad masses. Values are mass-adjusted least squares means and standard errors and accompanying p-values from SAS Procedure Mixed with body mass as a covariate. See Supplemental Table 1 for additional statistical details.





## Supplemental Material

Supplemental Table 1 is an Excel spreadsheet that summarizes p-values for main and interactive effects of linetype and treatment on various traits (most of which are reported in the manuscript). Subsequent Excel sheets contain more detailed results (degrees of freedom, least squares means, standard errors, etc.) for each traits, which can be found by navigating to the sheet corresponding to the “index” number indicated in column A of the RESULTS sheet.



Click here to access/download

**e-Component**

**G77\_SAS\_Results\_63\_UPLOAD.xlsx**

